For Research Use Only

CLASP1 Polyclonal antibody

Catalog Number: 20803-1-AP




Basic Information	Catalog Number: 20803-1-AP	GenBank Accession Numbe BC 112940	r: Purification Method: Antigen affinity purification
	Size: 300 µ g/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG14724	GeneID (NCBI): 23332 UNIPROT ID: Q7Z460 Full Name: cytoplasmic linker associat 1 Calculated MW: 1538 aa, 169 kDa Observed MW: 162 kDa	Recommended Dilutions: WB 1:500-1:3000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 ted protein
Applications	Tested Applications: WB,IP,IHC,ELISA Species Specificity: human, mouse, rat Note-IHC: suggested antige TE buffer pH 9.0; (*) Alterna retrieval may be performed buffer pH 6.0	WB SH-: IP:1 n retrieval with atively, antigen	itive Controls: : mouse brain tissue, HeLa cells, rat brain tissue, SY5Y cells HeLa cells, : human testis tissue,
Background Information	CLASP1 is a member of CLASP family. Members of the CLASP family of proteins are involved in multiple MT- dependent processes, including Golgi complex organization and trafficking, kinetochore fiber dynamics and flux, mitotic spindle assembly, and the stabilization of MT plus ends near the cortex. CLASP proteins regulate MT dynamics by promoting MT rescue and suppressing MT catastrophe. Recent work showed that CLASP1 activity is required for proper positioning of the mitotic spindle in both the C. elegans embryo and mammalian cells. Recent work has also identified roles CLASP1 in the regulation of spindle positioning.		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20803-1-AP (CLASP1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 20803-1-AP (CLASP1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 20803-1-AP (CLASP1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-CLASP1 (IP:20803-1-AP, 4ug; Detection:20803-1-AP 1:2000) with HeLa cells lysate 1750 ug.

Various lysates were subjected to SDS PAGE followed by western blot with 20803-1-AP (CLASP1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.