For Research Use Only

TCERG1L Polyclonal antibody

Isotype:

Catalog Number: 20858-1-AP



Purification Method:

IHC 1:50-1:500 IF-P 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 20858-1-AP BC042951 GeneID (NCBI): Size: 800 μ g/ml 256536 **UNIPROT ID:** Source: Rabbit

Q5VWI1 Full Name:

transcription elongation regulator 1-

Immunogen Catalog Number:

AG14828 Calculated MW: 586 aa, 66 kDa

Applications

Tested Applications: Positive Controls: IHC, IF-P, ELISA IHC: mouse testis tissue, Species Specificity: IF-P: mouse testis tissue, human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

TCERG1L, Transcription elongation regulator 1-like protein, is associated with plasma adiponectin, and plasma $adiponectin \ is \ a \ key \ modulator \ of \ obesity, in flammation, IR \ and \ diabetes \ (PMID: 22791750). \ Gene \ silencing \ of \ FBN2$ and TCERG1L is associated with promoter DNA hypermethylation in CRC tumors and may be biomarkers for the early detection of CRC (PMID:22238052). Stra8 and TCERG1L play a role in retinoic acid induced spermatogonial differentiation in mouse (PMID:3395975).

Storage

Storage:


Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data

Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 20858-1-AP (TCERG1L antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using TCERG1L antibody (20858-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 20858-1-AP (TCERG1L antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).