For Research Use Only

GABPB2 Polyclonal antibody Catalog Number: 20870-1-AP 1 Publications



Basic Information	Catalog Number: 20870-1-AP	GenBank Accession Number: BC027033	Purification Method: Antigen affinity purification	
	Size: 900 ug/ml	GenelD (NCBI): 126626	Recommended Dilutions: WB 1:500-1:1000	
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG14933	UNIPROT ID: Q8TAK5		
		Full Name: GA binding protein transcription factor, beta subunit 2		
		Calculated MW: 448 aa, 49 kDa		
		Observed MW: 49 kDa		
Applications	Tested Applications: WB, ELISA	Positive C	Positive Controls: WB : Jurkat cells, A375 cells, HEK-293 cells, HeLa cells	
	Cited Applications:	WB : Jurka		
	Species Specificity: human, mouse, rat			
	Cited Species: human			
Background Information	GABPB2 is a subunit of the GA-binding protein (GABP) transcription factor, which is a heterodimeric complex. The GABP complex plays a role in the regulation of gene transcription by binding to specific DNA sequences. GABPB2 specifically contributes to the transactivation of target genes (PMID: 18628204). The protein is primarily located in the nucleus. It is ubiquitously expressed in various tissues, including testis, ovary, and others. Studies have shown that GABPB2 is dispensable for normal lymphocyte development but moderately affects B cell responses (PMID: 18628204). Mice deficient in GABPB2 exhibit increased B cell proliferation and antibody production in response to certain stimuli. The molecular weight of GABPB2 is 48 kDa.			
Notable Publications	Author	Pubmed ID Journal	Application	
	Zhumei Shi	36270249 Mol Cell	IP	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide ar Aliquoting is unnecessary for -20	nd 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 20870-1-AP (GABPB2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.