For Research Use Only

HB9 Polyclonal antibody


Catalog Number:20975-1-AP



Basic Information	Catalog Number: 20975-1-AP	GenBank Accession Number: NM_005515	Purification Method: Antigen affinity purification	
	Size: 300 µg/ml	GenelD (NCBI): 3110	Recommended Dilutions: WB 1:500-1:1000	
	Source: Rabbit	UNIPROT ID: P50219		
	lsotype: IgG	Full Name: motor neuron and pancreas homeobox 1 Calculated MW: 41 kDa		
				Observed MW: 45-50 kDa
		Applications	Tested Applications: WB, ELISA Species Specificity: human, mouse	Positive Controls:
WB : mou	se spleen tissue, mouse liver tissue			
Background Informati	differentiation and pancreat MNX1 is associated with the with β cell maturation . MNX inhibit their ability to differe addition, the association bet Currarino syndrome phenoty	Motor neuron and pancreas homeobox (MNX1) is a homologous box transcription factor that promotes motor neuron differentiation and pancreatic development. MNX1 knockout in mice indicated that only the early expression of MNX1 is associated with the formation of pancreatic buds, and MNX1 is expressed in late β cells and associated with β cell maturation . MNX1 has the ability to induce endocrine progenitor cells to differentiate into β cells and inhibit their ability to differentiate into α cells, thereby maintaining the balance between α and β cells. In addition, the association between MNX1 gene mutations and Currarino syndrome has also been demonstrated. The Currarino syndrome phenotype caused by a MNX1 mutations was demonstrated to be a result of haploinsufficiency . A recent study also indicated that MNX1 may be an oncogene in prostate cancer (PMID: 30066929).		
Storage	Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azid Aliquoting is unnecessary fo	e and 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse spleen tissue were subjected to SDS PAGE followed by western blot with 20975-1-AP (HB9 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.