## For Research Use Only OTUD5 Polyclonal antibody Catalog Number: 21002-1-AP 8 Publications




Basic Information	Catalog Number: 21002-1-AP	L-AP BC028225 Antigen tration: GeneID (NCBI): Recomm ml 55593 WB 1:500 UNIPROT ID: IF/ICC 1: Q96G74		Purification Method: Antigen affinity purification
	Concentration: 600 ug/ml			Recommended Dilutions: WB 1:500-1:3000 IF/ICC 1:50-1:500
	Source: Rabbit			
	lsotype: IgG			
	Immunogen Catalog Number:Calculated MW:AG15190571 aa, 61 kDaObserved MW:70-75 kDa			
Applications	Tested Applications:	Positive Controls:		
	WB, IF/ICC, ELISA Cited Applications:			93T cells, HEK-293 cells, HeLa cells, Jurka 2 cells, NIH/3T3 cells, C6 cells
	WB, IF, IP, CoIP IF/ICC : A431 cells, Species Specificity: human, mouse, rat			
	Background Information	The deubiquitinating enzyme OTUD5, also known as deubiquitinating enzyme A (DUBA), is a member of the deubiquitinating enzyme OTU family. By deubiquitinating other proteins, OTUD5 is an important regulator of several physiological processes, such as immune signaling and DNA damage responses. The dysregulation of these processes can increase the risk of associated diseases, including inflammation, cancer and genetic disorders.		
Notable Publications	Author	Pubmed ID	Journal	Application
	Qiangqiang Liu	35913115	Elife	WB
	Junjing Zhang	40070026	Clin Transl Med	WB
	Xiaomeng Song	39721018	Adv Sci (Weinh)	WB,IF
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer:	ar after shipment.		


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



HEK-293T cells were subjected to SDS PAGE followed by western blot with 21002-1-AP (OTUD5 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 21002-1-AP (OTUD5 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using OTUD5 antibody (21002-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).