For Research Use Only

RNF213 Polyclonal antibody

Catalog Number: 21028-1-AP 1 Publications



Basic Information

Catalog Number:
21028-1-AP
BC032220
Concentration:
750 ug/ml
Source:
Rabbit
UNIPROT ID:
Rabce:
Full Name:

IgG ring finger protein 213
Immunogen Catalog Number: Calculated MW:
AG15335 3280 aa, 374 kDa

Purification Method: Antigen Affinity purified Recommended Dilutions: WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications: IHC

Species Specificity:

human Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HUVEC cells, IP: HUVEC cells,

IHC: human lymphoma tissue, human spleen tissue

Background Information

RNF213 is a typical E3 ubiquitin ligase that can catalyze ubiquitination of both proteins and lipids, and which is involved in various processes, such as lipid metabolism, angiogenesis and cell-autonomous immunity (PMID: 21799892, 26126547, 26278786, 26766444, 30705059).RNF213 contains a C3HC4-type RING finger domain, which is a specialized type of Zn-finger that binds two atoms of zinc and is thought to be involved in mediating protein-protein interactions.

Notable Publications

Author	Pubmed ID	Journal	Application
Ning Jiang	33200540	Thorac Cancer	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Immunohistochemical analysis of paraffinembedded human lymphoma slide using 21028-1-AP (RNF213 Antibody) at dilution of 1:50.



Immunohistochemical analysis of paraffinembedded human lymphoma slide using 21028-1-AP (RNF213 Antibody) at dilution of 1:50.



Various lysates were subjected to SDS PAGE followed by western blot with 21028-1-AP (RNF213 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-RNF213 (IP:21028-1-AP, 4ug; Detection:21028-1-AP 1:300) with HUVEC cells lysate 910 ug.