For Research Use Only

PML Polyclonal antibody Catalog Number: 21041-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: 21041-1-AP	GenBank Accession Number: BC020994	Purification Method: Antigen affinity purification
	Concentration:	GenelD (NCBI):	Recommended Dilutions:
	600 ug/ml	5371	WB 1:1000-1:4000
	Source:	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
	Rabbit	P29590	IHC 1:100-1:400
	Isotype: IgG	Full Name: promyelocytic leukemia	IF/ICC 1:50-1:500
	Immunogen Catalog Number: AG15377	Calculated MW: 882 aa, 98 kDa	
		Observed MW: 120 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, IF/ICC, IP, ELISA	WB : He	La cells, HEK-293 cells
	Cited Applications: WB, IHC, IF	P HEK-293 Cells.	
	Species Specificity:		
	human		
	Cited Species: human		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Information	PML, also named as MYL, RNF71 and TRIM19, is a member of RBCC/TRIM family of proteins that possess a RING finger domain, B-box, and coiled-coil domain. It functions as tumor suppressor. PML is required for normal, caspase-dependent apoptosis in response to DNA damage, FAS, TNF, or interferons. It plays a role in transcription regulation DNA damage response, DNA repair and chromatin organization. PML plays a role in processes regulated by retinoic acid, regulation of cell division, terminal differentiation of myeloid precursor cells and differentiation of neural progenitor cells. It regulates PTEN compartmentalization through the inhibition of USP7-mediated deubiquitinylation. A large number of alternative spliced transcripts are synthesized from the PML gene, resulting i a variety of PML proteins ranging in molecular weight from 48-97 kDa (PMID:11704850). And it can be detected as 70-130 kDa or larger due to the modification (especially SUMO) (PMID: 16778193, 22438555).		
	acid, regulation of cell division, to progenitor cells. It regulates PTEN deubiquitinylation. A large numb a variety of PML proteins ranging	l compartmentalization through the er of alternative spliced transcripts in molecular weight from 48-97 kD	inhibition of USP7-mediated are synthesized from the PML gene, resulting a (PMID:11704850). And it can be detected as
Notable Publications	acid, regulation of cell division, to progenitor cells. It regulates PTEN deubiquitinylation. A large numb a variety of PML proteins ranging 70-130 kDa or larger due to the m	I compartmentalization through the er of alternative spliced transcripts in molecular weight from 48-97 kD odification (especially SUMO) (PMI	inhibition of USP7-mediated are synthesized from the PML gene, resulting a (PMID:11704850). And it can be detected as D: 16778193, 22438555).
Notable Publications	acid, regulation of cell division, to progenitor cells. It regulates PTEN deubiquitinylation. A large numb a variety of PML proteins ranging 70-130 kDa or larger due to the m	I compartmentalization through the er of alternative spliced transcripts in molecular weight from 48-97 kD odification (especially SUMO) (PMI Pubmed ID Journal	inhibition of USP7-mediated are synthesized from the PML gene, resulting a (PMID:11704850). And it can be detected as D: 16778193, 22438555). Application
Notable Publications	acid, regulation of cell division, to progenitor cells. It regulates PTEN deubiquitinylation. A large numb a variety of PML proteins ranging 70-130 kDa or larger due to the m Author Orhi Barroso-Gomila	I compartmentalization through the er of alternative spliced transcripts in molecular weight from 48-97 kD odification (especially SUMO) (PMI Pubmed ID Journal 34795231 Nat Commun	inhibition of USP7-mediated are synthesized from the PML gene, resulting a (PMID:11704850). And it can be detected as D: 16778193, 22438555). Application WB,IF
Notable Publications	acid, regulation of cell division, to progenitor cells. It regulates PTEN deubiquitinylation. A large numb a variety of PML proteins ranging 70-130 kDa or larger due to the m Author Orhi Barroso-Gomila Jiaojiao Shen	I compartmentalization through the er of alternative spliced transcripts in molecular weight from 48-97 kD odification (especially SUMO) (PMI Pubmed ID Journal	inhibition of USP7-mediated are synthesized from the PML gene, resulting a (PMID:11704850). And it can be detected as D: 16778193, 22438555). Application

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of PML antibody (21041-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PML transfected HeLa cells.




HEK-293 cells were subjected to SDS PAGE followed by western blot with 21041-1-AP (PML Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




IP result of anti-PML (IP:21041-1-AP, 4ug; Detection:21041-1-AP 1:500) with HEK-293 cells lysate 3200ug.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 21041-1-AP (PML Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 21041-1-AP (PML Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using PML antibody (21041-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PML antibody (21041-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).