

For Research Use Only

NR1I3 Polyclonal antibody

Catalog Number: 21042-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number:

21042-1-AP

Size:

300 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG15378

GenBank Accession Number:

BC069651

GeneID (NCBI):

9970

UNIPROT ID:

Q14994

Full Name:

nuclear receptor subfamily 1, group I, member 3

Calculated MW:

352 aa, 40 kDa

Observed MW:

45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:250-1:1000

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HepG2 cells, L02 cells

IHC : mouse kidney tissue, rat liver tissue

Background Information

The nuclear hormone receptor superfamily is a large group of related transcription factors which includes members that bind a diverse array of ligands, including steroids, retinoic acid, thyroid hormone, and vitamin D. Nuclear hormone receptors contain a DNA-binding domain (DBD) and a ligand-binding domain [PMID: 10221899]. Nuclear receptor subfamily 1 group I member 3 (NR1I3; CAR), binds and transactivates the retinoic acid response elements that control expression of the retinoic acid receptor beta 2 and alcohol dehydrogenase 3 genes. It is expressed primarily in liver and regulates the expression of genes involved in xenobiotic metabolism as well as hormone, energy, and lipid homeostasis [PMID:22896671].

Notable Publications

Author	Pubmed ID	Journal	Application
Na Yao	31611795	Front Pharmacol	WB
Qingqing Hu	30414935	Biochem Pharmacol	WB
Han Li	33896369	Xenobiotica	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 21042-1-AP (NR1/3 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 21042-1-AP (NR1/3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 21042-1-AP (NR1/3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).