For Research Use Only

MLKL Polyclonal antibody

Catalog Number: 21066-1-AP

Featured Product

63 Publications



Basic Information

Catalog Number:

21066-1-AP

BC028141

Size:

450 µg/ml

197259

Source:

Rabbit

Q8NB16

Isotype:

GenBank Accession Number:

BC028141

GeneID (NCBI):

197259

UNIPROT ID:

Q8NB16

Full Name:

mixed lineage kinase domain-like

Immunogen Catalog Number:Calculated MW:AG15188471 aa, 54 kDaObserved MW:

54 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF, IP

human Cited Species: human, rat, pig

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HT-29 cells, HepG2 cells

IHC: human liver cancer tissue, human kidney tissue

Purification Method:

WB 1:500-1:3000 IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IF/ICC : HepG2 cells,

Background Information

Mixed lineage kinase domain like pseudokinase (MLKL), belongs to the protein kinase superfamily, has two MW of 54 and 30 kDa. MLKL plays a critical role in tumor necrosis factor (TNF)-induced necroptosis, a programmed cell death process, via interaction with receptor-interacting protein 3 (RIP3), which is a key signaling molecule in necroptosis pathway. High levels of this protein and RIP3 are associated with inflammatory bowel disease in children.

Notable Publications

Author	Pubmed ID	Journal	Application
Qing-Song Lin	31506635	Lab Invest	WB
Yueyang Wang	34480665	Biol Trace Elem Res	WB
Muhammad Khan	36389725	Front Immunol	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21066-1-AP (MLKL antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 21066-1-AP (MLKL antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 21066-1-AP (MLKL antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using MLKL antibody (21066-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP result of anti-MLKL (IP:21066-1-AP, 4ug; Detection:21066-1-AP 1:1000) with HepG2 cells lysate 1470 ug.