

NSF Polyclonal antibody

Catalog Number: 21172-1-AP

1 Publications

Basic Information

Catalog Number:

21172-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG15422

GenBank Accession Number:

BC030613

GeneID (NCBI):

4905

UNIPROT ID:

P46459

Full Name:

N-ethylmaleimide-sensitive factor

Calculated MW:

744 aa, 83 kDa

Observed MW:

83 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:100-1:400

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

IF

Species Specificity:

human, mouse, rat

Cited Species:

mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse brain tissue, mouse cerebellum tissue, rat brain tissue

IP : mouse brain tissue,

IHC : mouse brain tissue,

Background Information

The N-ethylmaleimide-sensitive fusion protein (NSF) is a 70-83 kDa cytosol protein that is expressed in all cell types. NSF functions as an ATPase and it is a part of the 20S membrane fusion apparatus with the vesicular SNAP receptor (v-SNARE) synaptobrevin, and the target SNAREs (t-SNARE) syntaxin and SNAP-25. With the hydrolysis of ATP by NSF, the released energy may set-up or drive membrane fusion processes. NSF is required for general membrane transport and trafficking.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhaokang Cui	35017645	Cell Death Differ	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21172-1-AP (NSF antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-NSF (IP:21172-1-AP, 4ug; Detection:21172-1-AP 1:300) with mouse brain tissue lysate 3000ug.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 21172-1-AP (NSF Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieved with Sodium Citrate buffer (pH 6.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 21172-1-AP (NSF Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieved with Sodium Citrate buffer (pH 6.0).