For Research Use Only

PAX3 Polyclonal antibody

Catalog Number: 21386-1-AP 3 Publications



Basic Information

Catalog Number: 21386-1-AP

BC101301

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

GenBank Accession Number:

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

700 μg/ml Source:

5077 **UNIPROT ID:** WB 1:500-1:3000

Rabbit Isotype:

AG16021

P23760 Full Name:

paired box 3

Calculated MW: Immunogen Catalog Number:

484 aa, 53 kDa

Observed MW: 56-60 kDa

protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

WB: mouse skin tissue, HEK-293 cells

WB, IHC

Species Specificity:

human, mouse, rat

human, mouse, hamster

Cited Species:

IP: mouse skin tissue,

Positive Controls:

Background Information

PAX3, a transcription factor and multifunctional regulatory protein, is normally expressed during embryonic development. In the nervous system, PAX3 is involved in neural tube closure, neural crest development, and peripheral neuron differentiation. In the present study, PAX3 was reported as a novel regulator of GFAP transcription, and the overexpression and suppression of PAX3 could inhibit and promote NSC differentiation, respectively. In muscle development, PAX3 ensures the survival of myogenic progenitor cells with Pax3-expressing progenitors giving rise to both skeletal and smooth muscle cells. PAX3 also has a well-established role in the development of melanocytes during embryogenesis, and has recently been characterized as a molecular switch in the mature melanocyte. Mutations in PAX3 can cause Waardenburg syndrome.

Notable Publications

Author	Pubmed ID	Journal	Application
Chuanhui Yin	32949110	Birth Defects Res	IHC
Meng-Han Zhang	39758702	Stem Cells Int	WB
Li Zhang	39307459	Neurochem Int	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



mouse skin tissue were subjected to SDS PAGE followed by western blot with 21386-1-AP (PAX3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-PAX3 (IP:21386-1-AP, 4ug; Detection:21386-1-AP 1:500) with mouse skin tissue lysate 3200ug.