For Research Use Only

HIGD2A Polyclonal antibody

Catalog Number: 21415-1-AP

Featured Product

2 Publications



Basic Information

Applications

Catalog Number: 21415-1-AP Size: 300 μg/ml Source: Rabbit

Isotype:

Immunogen Catalog Number:

AG15892

Tested Applications: WB, IF/ICC, ELISA

Cited Applications:

Species Specificity: human, mouse Cited Species: human

GenBank Accession Number:

BC007502 GeneID (NCBI): 192286 **UNIPROT ID:** Q9BW72 Full Name:

HIG1 hypoxia inducible domain family, member 2A

Calculated MW: 106 aa, 12 kDa Observed MW: 11 kDa, 50 kDa

Antigen affinity purification

Purification Method:

Recommended Dilutions: WB 1:500-1:1000 IF/ICC 1:200-1:800

Positive Controls: WB: mouse kidney tissue,

IF/ICC: MCF-7 cells,

Background Information

The mammalian HIGD-family proteins include two subgroups of yeast Rcf1 homologs, HIGD1A/B/C and HIGD2A/B. HIGD1A and HIGD2A (of 10.4 and 11.5 kDa, respectively) have two hydrophilic transmembrane domains and are localized in the inner mitochondrial membrane. HIGD1A and HIGD2A expression is induced under stress conditions like hypoxia or glucose deprivation in human cervical epithelial cells and murine cerebral cortical neurons. HIGD1A may play pleiotropic roles in mitochondrial respiratory chain biogenesis. HIGD2A forms a 50 kDa regulatory module with COX4-1, COX5B, and COX6A1, which binds to newly synthesized COX3 to promote its binding to the previously assembled COX1 + COX2 module (PMID:33291261,32375044).

Notable Publications

Author	Pubmed ID	Journal	Application
Daniella H Hock	33451405	Mol Cell Proteomics	WB
Daniella H Hock	32317297	Mol Cell Proteomics	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 21415-1-AP (HIGD2A antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using HIGD2A antibody (21415-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).