For Research Use Only

VANGL2 Polyclonal antibody

Catalog Number: 21492-1-AP

Featured Product

10 Publications



Basic Information

Catalog Number: GenBank Accession Number: 21492-1-AP BC103920 GeneID (NCBI): Size: 450 µg/ml 57216 **UNIPROT ID:** Source: Rabbit Q9ULK5 Full Name: Isotype:

vang-like 2 (van gogh, Drosophila)

Calculated MW: Immunogen Catalog Number: AG15833 521 aa, 60 kDa Observed MW: 60-70 kDa

Tested Applications:

Cited Applications: WB, IP, IF, CoIP

human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:1000

WB: HepG2 cells, mouse brain tissue, NIH/3T3 cells,

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF 1:20-1:200

Positive Controls:

rat brain tissue

IP: HepG2 cells,

IF: HepG2 cells,

IHC: human stomach tissue,

Applications

IF/ICC, IHC, IP, WB, ELISA

Species Specificity:

human, rat, mouse

Background Information

Vangl2 is a key component of the planar cell polarity (PCP) pathway. Vangl2 is tightly associated with the postsynaptic density (PSD) fraction and forms a protein complex with PSD-95 and NMDA receptors. Vangl2 directly binds to the third PDZ domain of PSD-95 via its C-terminal TSV motif. Vangl2 directly binds to N-cadherin.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiao Han	36054333	Hum Mutat	WB
Tammy N Jessen	29097183	Exp Cell Res	CoIP
Xin Sheng	34738156	Cell Tissue Res	WB,IF,IP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21492-1-AP (VANGL2 antibody) at dilution of 1:800 incubated at room temporarium for 1.5 hours




Various lysates were subjected to SDS PAGE followed by western blot with 21492-1-AP (VANGL2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




IP result of anti-VANGL2 (IP:21492-1-AP, 4ug; Detection:21492-1-AP 1:500) with HepG2 cells lysate 1880 ug.



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 21492-1-AP (VANGL2 antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 21492-1-AP (VANGL2 antibody) at dilution of 1:200 (under 40x lens).



Immun of luorescent analysis of HepG2 cells using 21492-1-AP (VANGL2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Affini Pure Goat Anti-Rabbit IgG(H+L).