#### For Research Use Only

# GABPA Polyclonal antibody

Catalog Number: 21542-1-AP

Featured Product

14 Publications

BC035031

GenBank Accession Number:



**Basic Information** 

Catalog Number: 21542-1-AP

Concentration: GeneID (NCBI): 500 µ g/ml 2551

Source: UNIPROT ID:
Rabbit Q06546
Isotype: Full Name:

IgG GA binding protein transcription

Immunogen Catalog Number: factor, alpha subunit 60kDa

AG16191 Calculated MW: 454 aa, 51 kDa

Observed MW: 56-60 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:200-1:800

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, ChIP Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293 cells, A431 cells, mouse liver tissue, MCF-7 cells, A549 cells, rat brain tissue, K-562 cells, NIH/3T3 cells, mouse brain tissue

IP: HeLa cells.

IHC: human cervical cancer tissue, human breast

cancer tissue

IF/ICC: HEK-293 cells,

## **Background Information**

GA-binding protein alpha chain (GABP alpha subunit, GABPA, nuclear respiratory factor 2 subunit alpha, transcription factor E4TF1-60) is one of three GA-binding protein transcription factor subunits which functions as a DNA-binding subunit.GABPA is a member of Ets family, binds to the Yap promoter and activates YAP transcription(23684612). Since this subunit shares identity with a subunit encoding the nuclear respiratory factor 2 gene, it is likely involved in activation of cytochrome oxidase expression and nuclear control of mitochondrial function. This subunit also shares identity with a subunit constituting the transcription factor E4TF1, responsible for expression of the adenovirus E4 gene. Because of its chromosomal localization and ability to form heterodimers with other polypeptides, this gene may play a role in the Down Syndrome phenotype.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Narendra Kumar Verma	28905448	Stem Cells	WB
Shaofan Hu	36174386	Redox Biol	WB
Sheng Zhang	28549418	BMC Cancer	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 21542-1-AP (GABPA antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer using 21542-1-AP (GABPA antibody) at dilution of 1:50 (under 10x lens)




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 21542-1-AP (GABPA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 21542-1-AP (GABPA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-GABPA (IP:21542-1-AP, 4ug; Detection:21542-1-AP 1:1000) with HeLa cells lysate 1200ug.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using GABPA antibody (21542-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).