

For Research Use Only

# NAGS Polyclonal antibody

Catalog Number: 21566-1-AP **1 Publications**



## Basic Information

<b>Catalog Number:</b> 21566-1-AP	<b>GenBank Accession Number:</b> BC111713	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 133 µg/ml	<b>GeneID (NCBI):</b> 162417	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IHC 1:20-1:200 IF 1:10-1:100
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q8N159	
<b>Isotype:</b> IgG	<b>Full Name:</b> N-acetylglutamate synthase	
<b>Immunogen Catalog Number:</b> AG16073	<b>Calculated MW:</b> 534 aa, 58 kDa	
	<b>Observed MW:</b> 58-66 kDa	

## Applications

<b>Tested Applications:</b> IF/ICC, IHC, WB, ELISA	<b>Positive Controls:</b> WB : HepG2 cells, Jurkat cells, LO2 cells IHC : human liver tissue, human small intestine tissue IF : HepG2 cells, Hela cells
<b>Cited Applications:</b> WB	
<b>Species Specificity:</b> human, mouse, rat	
<b>Cited Species:</b> mouse	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

N-acetylglutamate synthase, also known as NAGS, is a mitochondrial enzyme that catalyzes the formation of N-acetylglutamate (NAG) from glutamate and acetyl coenzyme-A. NAGS is the key enzyme for regulating the hepatic urea cycle and is also highly expressed in the kidney and gut. Deficiency of NAGS leads to hyperammonemia due to decreased activity of CPS1 deprived of its cofactor NAG(PMID:12459178,12754705).

## Notable Publications

Author	Pubmed ID	Journal	Application
Wenxiang Zhang	32029232	Mol Metab	WB

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 21566-1-AP (NAGS antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver using 21566-1-AP (NAGS antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver using 21566-1-AP (NAGS antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells, using NAGS antibody 21566-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).