For Research Use Only

NRCAM Polyclonal antibody

Catalog Number:21608-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number:

21608-1-AP

Size:

800 µg/ml

Source:

Rabbit

GenBank Accession Number:

BC098401

GeneID (NCBI):

4897

UNIPROT ID:

Q92823

Isotype:

Full Name:

neuronal cell adhesion molecule

Immunogen Catalog Number:Calculated MW:AG162371304 aa, 144 kDa

Observed MW: 130-136 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: mouse brain tissue, human brain tissue, HEK-293

Purification Method:

WB 1:500-1:2000 IHC 1:200-1:800

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

cells, rat brain tissue

IHC: mouse brain tissue, human brain tissue

IF/ICC: HEK-293 cells,

Background Information

Neuronal cell adhesion molecule (NRCAM) is a member of the L1 subfamily of cell adhesion molecules (CAMs) that belong to the immunoglobulin superfamily (PMID: 11329126). NRCAM is a transmembrane protein composed of six Ig-like domains and five fibronectin type-III repeats in the extracellular region, with a highly conserved cytoplasmic tail. It is mainly expressed in the nervous system and is involved in neuron-neuron adhesion and promotes directional signaling during axonal cone growth. NRCAM is also expressed outside the nervous system. Altered expression of NRCAM has been associated with tumor progression in diverse organs (PMID: 22182708).

Notable Publications

Author	Pubmed ID	Journal	Application
Zhen-Huang Chen	32918999	Mol Cell Neurosci	IHC
Michela Lupia	29551673	Stem Cell Reports	IF
Hongwei Jin	34150622	Front Oncol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21608-1-AP (NRCAM antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 21608-1-AP (NRCAM antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 21608-1-AP (NRCAM antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using 21608-1-AP (NRCAM antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).