## For Research Use Only

## STRN Polyclonal antibody

Catalog Number:21624-1-AP

Featured Product

**6 Publications** 



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 21624-1-AP
 BC 106879

 Size:
 GeneID (NCBI):

 240 µg/ml
 6801

 Source:
 UNIPROT ID:

Rabbit O43815
Isotype: Full Name:

gG striatin, calmodulin binding protein

Immunogen Catalog Number:Calculated MW:AG15984780 aa, 86 kDaObserved MW:

110 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls

WB: A549 cells, rat brain tissue, NIH/3T3 cells, mouse brain tissue, HEK-293T cells, HeLa cells, Jurkat cells

**Purification Method:** 

WB 1:2000-1:10000

protein lysate

IHC 1:50-1:500

IF/ICC 1:10-1:100

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP: mouse brain tissue,

IHC: mouse brain tissue, human liver tissue

IF/ICC: Hela cells,

**Background Information** 

STRN (Striatin), a 780-amino acid protein, was identified and cloned more than a decade ago. In adult mammals, striatin is mainly expressed in neurons of both the central and peripheral nervous systems with very high levels of expression in the striatum, hence its name. STRN-ALK fusion may be a potential therapeutic target in the aggressive forms of thyroid cancer (PMID: 24475247).

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Daniel Meyer	36292701	Genes (Basel)	WB
Qian-Yu Du	32280692	Biomed Res Int	WB,IHC
Liping Chen	24962574	J Biol Chem	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 21624-1-AP (STRN antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




IP result of anti-STRN (IP:21624-1-AP, 4ug; Detection:21624-1-AP 1:4000) with mouse brain tissue lysate 1080 ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 21624-1-AP (STRN antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of Hela cells, using STRN antibody 21624-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human STRN (21624-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit I gG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).