For Research Use Only

IDE Polyclonal antibody

Catalog Number: 21728-1-AP

Featured Product

11 Publications



Basic Information

Catalog Number:

21728-1-AP

BC096336

Size:

GeneID (NCBI):

300 µg/ml

3416

Source:

UNIPROT ID:

Rabbit

P14735

Isotype:

GenBank Accession Number:

BC096336

UNCBI):

3416

UNIPROT ID:

Full Name:

gG Abeta degrading protease

Immunogen Catalog Number: Calculated MW: AG16392 1019 aa, 118 kDa

Observed MW: 100-120 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB, IHC

Species Specificity: human

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A2780 cells, HepG2 cells, HeLa cells, DU 145 cells

IP: HepG2 cells,

IHC: human kidney tissue, human liver tissue

Background Information

IDE (INS-degrading enzyme) belongs to the peptidase M16 family. It is also known as insulysin and is a 110-kD neutral metallopeptidase that can degrade a number of peptides, including INS and beta-amyloid(PMID:9830016). The reduction of IDE expression may be a risk factor for AD, and that IDE may interact with APOE status to affect beta-amyloid metabolism(PMID:12507914). It can exsit as a homodimer and form higher oligomers(PMID:17613531). This antibody is specific to IDE.

Notable Publications

Author	Pubmed ID	Journal	Application
Ke-Wei Chang	30217465	Neurochem Int	WB
Chunling Yuan	30389579	Biochim Biophys Acta Mol Basis Dis	WB
Hui Zeng	32473218	Int J Biol Macromol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21728-1-AP (IDE antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human kidney using 21728-1-AP (IDE antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney using 21728-1-AP (IDE antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-IDE (IP:21728-1-AP, 4ug; Detection:21728-1-AP 1:500) with HepG2 cells lysate 1600ug.



WB result of IDE antibody (21728-1-AP; 1:400; incubated at room temperature for 1.5 hours) with sh-Control and sh-IDE transfected HeLa cells.