For Research Use Only

KANK2 Polyclonal antibody

Catalog Number:21733-1-AP 4 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

| Basic Information | Catalog Number: 21733-1-AP | GenBank Accession Num BC098105 | | Method: nity purification | |
|------------------------|--|-----------------------------------|--------------------------|--|--|
| | Size: | GenelD (NCBI): | Recommend | led Dilutions: | |
| | 500 µg/ml | 25959 | WB 1:500-1: | | |
| | Source: Rabbit | UNIPROT ID: Q63ZY3 | | IHC 1:50-1:500 IF/ICC 1:50-1:500 | |
| | Isotype: Full Name: IgG KN motif and ankyrin repeat domains | | | | |
| | Immunogen Catalog Number: AG16416 | 2 | | | |
| | | | | | |
| | | Observed MW: 95-100 kDa | | | |
| Applications | Tested Applications: | Positive Controls: | | | |
| | IF/ICC, IHC, WB, ELISA Cited Applications: WB, IF | | | nouse heart tissue, mouse kidney tissue, ected HEK-293 cells, mouse lung tissue | |
| | Species Specificity: | I | C : human kidney tissue, | | |
| | human, mouse, rat | IF/ICC : HepG2 cells, A549 cells | | | |
| | Cited Species: human, mouse | | | | |
| | Note-IHC: suggested antig TE buffer pH 9.0; (*) Alter retrieval may be performe buffer pH 6.0 | natively, antigen | | | |
| Background Information | Involved in transcription regulation by sequestering nuclear receptor coactivators, such as NCOA1, NCOA2 and NCOA3, in the cytoplasm; the function is deregulated by phosphorylation. May be involved in the control of cytoskeleton formation by regulating actin polymerization. Involved in regulation of caspase-independent apoptosis; proposed to sequester AIFM1 in mitochondria and apoptotic stimuli lead to its proteasomal degradatio allowing the release of AIFM1 to the nucleus. May be involved in promotion of cell proliferation. | | | | |
| Notable Publications | Author | Pubmed ID Journal | L | Application | |
| | Pantipa Tonsawan | 31553647 Am J Pł | nysiol Cell Physiol | WB | |
| | Yuqun Xu | 34274317 J Biol C | hem | WB | |
| | Li Liu | 37927241 Am J Ph | nysiol Cell Physiol | WB | |
| Storage | Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide ar | | | | |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21733-1-AP (KANK2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 21733-1-AP (KANK2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 21733-1-AP (KANK2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using KANK2 antibody (21733-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using KANK2 antibody (21733-1-AP) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), Coralite®594 B23/NPM1 antibody (CL594-60096, Clone: 4F 12A3, red).



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using KANK2 antibody (21733-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).