

# Ataxin 2 Polyclonal antibody

Catalog Number: 21776-1-AP

Featured Product

33 Publications

## Basic Information

## Catalog Number:

21776-1-AP

## Size:

350 µg/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG16470

## GenBank Accession Number:

BC114546

## GeneID (NCBI):

6311

## UNIPROT ID:

Q99700

## Full Name:

ataxin 2

## Calculated MW:

1313 aa, 140 kDa

## Observed MW:

140-150 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:2000-1:16000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF-P 1:50-1:500

IF/ICC 1:450-1:1800

## Applications

## Tested Applications:

WB, IHC, IF/ICC, IF-P, IP, ELISA

## Cited Applications:

WB, IHC, IF, IP, CoIP

## Species Specificity:

human, mouse, rat

## Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

WB : HeLa cells, HEK-293 cells, Jurkat cells, Neuro-2a cells

IP : HEK-293 cells,

IHC : mouse brain tissue,

IF-P : mouse brain tissue,

IF/ICC : HepG2 cells,

## Background Information

ATXN2 contains a repeat structure with 22 or 23 triplets coding for glutamine and the (CAG)<sub>8</sub>CAA(CAG)<sub>4</sub>CAA(CAG)<sub>8</sub> sequence; expansion of this domain to a size ≥34 triplets with a pure CAG sequence primarily causes autosomal dominant SCA2 [PMID:18418684], while ATXN2 expansions with CAA interruptions were observed as the cause of Levo-dopa responsive Parkinson's disease [PMID:10668726]. ATXN2 expansions associated with ALS were reported to be interrupted by at least one CAA triplet [PMID:21537950]

## Notable Publications

Author	Pubmed ID	Journal	Application
Garam Kim	36288714	Cell Rep	WB,IF
Lauren A Ostrowski	30417124	Commun Biol	WB
Gaston Bonenfant	30944179	J Virol	WB,IF

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21776-1-AP (Ataxin 2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 21776-1-AP (Ataxin 2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Ataxin 2 antibody (21776-1-AP) at dilution of 1:900 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 21776-1-AP (Ataxin 2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Ataxin 2 (IP:21776-1-AP, 4ug; Detection:21776-1-AP 1:6000) with HEK-293 cells lysate 1800 ug.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Ataxin 2 antibody (21776-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).