

For Research Use Only

DSG2 Polyclonal antibody

Catalog Number: 21880-1-AP

Featured Product

8 Publications



Basic Information

Catalog Number:

21880-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG16393

GenBank Accession Number:

BC099655

GeneID (NCBI):

1829

UNIPROT ID:

Q14126

Full Name:

desmoglein 2

Calculated MW:

1118 aa, 122 kDa

Observed MW:

145-150 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:500-1:2000

IF 1:200-1:800

Applications

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

IF, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, HEK-293 cells, A431 cells, A549 cells, HeLa cells

IP: HepG2 cells,

IHC: human lung cancer tissue,

IF: A431 cells,

Background Information

Desmosomes are cell-cell junctions between epithelial, myocardial, and certain other cell types. Desmosomal cadherins, consisting of four desmogleins (DSG1-4) and three desmocollins (DSC1-3) in humans, mediate adhesion through calcium-dependent homophilic/heterophilic interactions. DSG2 is a single-pass transmembrane glycoprotein that is widely expressed in epithelial and non-epithelial tissues, such as the intestine, epidermis, testis, and heart (PMID:21715983). Defects in DSG2 are the cause of familial arrhythmogenic right ventricular dysplasia type 10 (ARVD10), and genetic variations in DSG2 are the cause of susceptibility to cardiomyopathy dilated type 1BB (CMD1BB).

Notable Publications

Author	Pubmed ID	Journal	Application
Jianhai Zhang	27237588	Food Chem Toxicol	WB
Chen Zhang	24826891	Helicobacter	IF
Indrajyoti Indra	34260281	Mol Biol Cell	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of DSG2 antibody (21880-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-DSG2 transfected HEK-293 cells.




HepG2 cells were subjected to SDS PAGE followed by western blot with 21880-1-AP (DSG2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




IP result of anti-DSG2 (IP:21880-1-AP, 4ug; Detection:21880-1-AP 1:1000) with HepG2 cells lysate 3600ug.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 21880-1-AP (DSG2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 21880-1-AP (DSG2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed A431 cells using DSG2 antibody (21880-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).