For Research Use Only

Sclerostin Polyclonal antibody

Catalog Number: 21933-1-AP

Featured Product

9 Publications



Basic Information

Catalog Number: 21933-1-AP

Size: 550 μg/ml

Source: Rabbit

Isotype:

BC101086 GeneID (NCBI): 50964

UNIPROT ID: Q9BQB4 Full Name: sclerosteosis

GenBank Accession Number:

Immunogen Catalog Number: AG16567

Calculated MW: 213 aa, 24 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IHC 1:50-1:500 IF 1:50-1:500

Applications

Tested Applications: IF/ICC, IHC, ELISA Cited Applications: WB, IF, IHC

Species Specificity: human, mouse **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human kidney tissue, human breast cancer tissue, mouse lung tissue, mouse embryo tissue

IF: HepG2 cells,

Background Information

Sclerostin is a secreted glycoprotein with a C-terminal cysteine knot-like (CTCK) domain and sequence similarity to the DAN (differential screening-selected gene aberrative in neuroblastoma) family of bone morphogenetic protein (BMP) antagonists. Loss-of-function mutations in this gene are associated with an autosomal-recessive disorder, sclerosteosis, which causes progressive bone overgrowth. A deletion downstream of this gene, which causes reduced sclerostin expression, is associated with a milder form of the disorder called van Buchem disease

Notable Publications

Author	Pubmed ID	Journal	Application
Fang He	33159018	Aging (Albany NY)	WB
Xiaoan Wei	35611810	EMBO Mol Med	IF
Marko Lucijanic	29532161	Ann Hematol	IHC

Storage


Storage:


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data







Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 21933-1-AP (Sclerostin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Sclerostin antibody (21933-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).