

For Research Use Only

FTCD Polyclonal antibody

Catalog Number: 21959-1-AP

5 Publications



Basic Information

Catalog Number:

21959-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG16646

GenBank Accession Number:

BC136395

GeneID (NCBI):

10841

UNIPROT ID:

O95954

Full Name:

formiminotransferase
cyclodeaminase

Calculated MW:

541 aa, 59 kDa

Observed MW:

53-62 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:4000-1:16000

IHC 1:250-1:1000

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : mouse liver tissue, rat liver tissue

IHC : human liver tissue,

Background Information

FTCD (formimidoyltransferase cyclodeaminase) is critical for the catabolism of histidine, a . This process is one-carbon units that can enter the one-carbon/folate cycle, which provides methyl groups for arsenic metabolism (PMID: 30893314). In addition, FTCD could serve as a predictive and prognostic marker for patients with HCC (PMID: 37675273).

Notable Publications

Author	Pubmed ID	Journal	Application
Shizhe Yu	35813194	Front Cell Dev Biol	IF
Jiajia Chen	31203308	Med Sci Monit	WB,IHC
Huimin Shen	33380233	Bioengineered	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21959-1-AP (FTCD antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 21959-1-AP (FTCD antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 21959-1-AP (FTCD antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).