For Research Use Only

N-cadherin Polyclonal antibody

Catalog Number:22018-1-AP

Featured Product

1295 Publications



Basic Information	22018-1-AP	atalog Number: GenBank Accession Number: 2018-1-AP BC036470		Purification Method: Antigen affinity purification					
	Concentration:	GeneID (NCBI): 1000 UNIPROT ID: P19022 Full Name: cadherin 2, type 1, N-cadherin (neuronal) Calculated MW: 906 aa, 100 kDa Observed MW:		Recommended Dilutions: WB 1:2000-1:16000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:2000-1:8000 IF-P 1:50-1:500 IF/ICC 1:200-1:800					
	650 ug/ml								
	Source: Rabbit								
	Isotype: IgG Immunogen Catalog Number: AG16792								
					130 kDa				
					Applications	Tested Applications: Positive Controls:		ntrols:	
						WB, IHC, IF/ICC, IF-P, IP, ELISA		WB: mouse brain tissue, A549 cells, C2C12 cells, HE	
		Cited Applications: WB, IHC, IF, CoIP				293 cells, rat brain tissue, HepG2 cells, NCI-H1299 cells, HeLa cells, C6 cells, PC-3 cells, rat heart tissue			
Species Specificity:		IP : mouse	IP : mouse brain tissue,						
human, mouse, rat		IHC : mous	IHC : mouse heart tissue, mouse brain tissue						
numan, mouse, rat, canine, bovine, namster, norse,		IF-P: mous	IF-P : mouse heart tissue, C2C12 cells						
		IF/ICC : C2	IF/ICC : C2C12 cells,						
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0									
		a with citrate							
Background Information	buffer pH 6.0 Cadherins are a family of transmiplay an important role in the mai as CDH2 (cadherin 2), is a 130-kD superfamily which also include E cell types including neurons, end cadherin has functions in early br	embrane glycoprotein ntenance of normal tis a transmembrane glyo -, P-, R-, and B-cadherin othelial cells and card ain morphogenesis, sy	ssue architecture. coprotein and a cl ns. Expression of l liac myocytes (PM ynaptogenesis an	assical member of the cadherin N-cadherin has been reported on various ID: 11282032; 9508779; 8125202). N- d synaptic plasticity (PMID: 23321619). Th					
	buffer pH 6.0 Cadherins are a family of transm play an important role in the mai as CDH2 (cadherin 2), is a 130-kD superfamily which also include E cell types including neurons, end cadherin has functions in early br N-cadherin ectodomain can be cle	embrane glycoprotein ntenance of normal tis a transmembrane glyc -, P-, R-, and B-cadherin othelial cells and card ain morphogenesis, sy eaved, leading to the g	ssue architecture. coprotein and a cl ns. Expression of l liac myocytes (PM ynaptogenesis an	N-cadherin (neural cadherin), also known assical member of the cadherin N-cadherin has been reported on various ID: 11282032; 9508779; 8125202). N- d synaptic plasticity (PMID: 23321619). Th kD N-terminal fragment (PMID: 1699883;					
	buffer pH 6.0 Cadherins are a family of transmiplay an important role in the mai as CDH2 (cadherin 2), is a 130-kD superfamily which also include E cell types including neurons, end cadherin has functions in early br N-cadherin ectodomain can be cle 17028923).	embrane glycoprotein ntenance of normal tis a transmembrane glyc -, P-, R-, and B-cadherin othelial cells and card ain morphogenesis, sy eaved, leading to the g	ssue architecture. coprotein and a cl ns. Expression of l liac myocytes (PM ynaptogenesis an generation of a 90	N-cadherin (neural cadherin), also known assical member of the cadherin N-cadherin has been reported on various ID: 11282032; 9508779; 8125202). N- d synaptic plasticity (PMID: 23321619). Th					
Background Information	buffer pH 6.0 Cadherins are a family of transmiplay an important role in the mai as CDH2 (cadherin 2), is a 130-kD superfamily which also include E cell types including neurons, end cadherin has functions in early br N-cadherin ectodomain can be cle 17028923).	embrane glycoprotein ntenance of normal tis a transmembrane glyc r, P-, R-, and B-cadherin othelial cells and card ain morphogenesis, sy eaved, leading to the g Pubmed ID J 36230734 (ssue architecture. coprotein and a cl ns. Expression of i liac myocytes (PM ynaptogenesis an generation of a 90	N-cadherin (neural cadherin), also known assical member of the cadherin N-cadherin has been reported on various ID: 11282032; 9508779; 8125202). N- d synaptic plasticity (PMID: 23321619). Th kD N-terminal fragment (PMID: 1699883) Application					
	buffer pH 6.0 Cadherins are a family of transmiplay an important role in the mai as CDH2 (cadherin 2), is a 130-kD superfamily which also include E cell types including neurons, end cadherin has functions in early br N-cadherin ectodomain can be cle 17028923).	embrane glycoprotein ntenance of normal tis a transmembrane glyc -, P-, R-, and B-cadherin othelial cells and card ain morphogenesis, sy eaved, leading to the g Pubmed ID J 36230734 (34638929	ssue architecture. coprotein and a cl ns. Expression of I liac myocytes (PM ynaptogenesis an generation of a 90 Journal Cancers (Basel)	N-cadherin (neural cadherin), also known assical member of the cadherin V-cadherin has been reported on various ID: 11282032; 9508779; 8125202). N- d synaptic plasticity (PMID: 23321619). Th kD N-terminal fragment (PMID: 1699883; Application WB					


W: ptgcn.com

T: 4006900926


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of N-cadherin antibody (22018-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-N-cadherin transfected HEK-293 cells.




C2C12 (N-cadherin+), HeLa (N-cadherin+), MCF-7 (N-cadherin-), A431 (N-cadherin-) cell lysates were subjected to SDS PAGE followed by western blot with 22018-1-AP (N-cadherin antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 22018-1-AP (Ncadherin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




IP result of anti-N-cadherin (IP:22018-1-AP, 3ug; Detection:22018-1-AP 1:1500) with mouse brain tissue lysate 7000ug.





Immunofluorescent analysis of (4% PFA) fixed mouse heart tissue using N-cadherin antibody (22018-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunofluorescent analysis of (4% PFA) fixed mouse heart tissue using N-cadherin antibody (22018-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).







Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 22018-1-AP (N-cadherin antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed C2C12 cells using N-cadherin antibody (22018-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).