For Research Use Only

iPLA2 Polyclonal antibody

Catalog Number:22030-1-AP

6 Publications



Purification Method:

WB 1:500-1:2000

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number:

22030-1-AP

BC036742

Size:

GeneID (NCBI):

650 µg/ml

8398

Source:

UNIPROT ID:

Rabbit

O60733

Isotype:

GenBank Accession Number:

BC036742

UNCBI):

8398

UNIPROT ID:

Full Name:

Full Name: phospholipase A2, group VI (cytosolic,

Immunogen Catalog Number: calcium-independent)
AG16896 Calculated MW:

90 kDa Observed MW: 80 kDa, 88 kDa

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB, IF, IHC Species Specificity: human, mouse

Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: mouse testis tissue,
IP: mouse testis tissue,

IHC: human small intestine tissue, human brain tissue

Background Information

PLA2G6, also named as PLPLA9, is involved in the remodeling of membranes allowing arachidonic acid to be placed in the proper position in phospholipids for stimulated release. It has been implicated in normal phospholipid remodeling, nitric oxide-induced or vasopressin-induced arachidonic acid release and in leukotriene and prostaglandin production. This protein has 4 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Yao Lu	31506141	Zhongguo Dang Dai Er Ke Za Zhi	IHC
Shuhua Shan	32657580	J Agric Food Chem	WB,IHC
Laura Mahoney-Sanchez	36001957	Cell Rep	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 22030-1-AP (iPLA2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-iPLA2 (IP:22030-1-AP, 4ug; Detection:22030-1-AP 1:800) with mouse testis tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 22030-1-AP (iPLA2 antibody at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 22030-1-AP (iPLA2 antibody at dilution of 1:50 (under 40x lens).