

For Research Use Only

FOXP1 Polyclonal antibody

Catalog Number: 22051-1-AP

2 Publications



Basic Information

Catalog Number:

22051-1-AP

Size:

800 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG17045

GenBank Accession Number:

BC131720

GeneID (NCBI):

27086

UNIPROT ID:

Q9H334

Full Name:

forkhead box P1

Calculated MW:

677 aa, 75 kDa

Observed MW:

50 kDa, 60-65 kDa, 85 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:250-1:1000

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, A549 cells, LNCaP cells, MCF-7 cells, MDA-MB-453s cells, mouse heart tissue, mouse lung tissue, mouse testis tissue, PC-3 cells, Raji cells, Y79 cells

IHC: mouse embryo tissue,

Background Information

FOXP1, also known as Mac-1-regulated forkhead, is a 677 amino acid protein, which forms homodimers and heterodimers with FOXP2 and FOXP4 (PubMed:25027557). Dimerization is required for DNA-binding. FOXP1 has an important function in neuronal development.⁹ Mutations of its gene, FOXP1, located on chromosome 3p14.1, can result in the development of autism spectrum disorder, intellectual disability, speech and language deficits as well as motor development delay. FOXP1 is also engaged in lung and esophagus morphogenesis, as well as in B-cell development.^{7,10} The widely researched role of FOXP1 in carcinogenesis is of great importance, although still unclear to some extent. FOXP1 exists some isoforms with MV 75-77 kDa, 65-67 kDa, 12 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Liangyi Cui	33372387	J Cell Mol Med	WB
Yunliang Liu	38861915	Int Immunopharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 22051-1-AP (FOXP1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse embryo tissue slide using 22051-1-AP (FOXP1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse embryo tissue slide using 22051-1-AP (FOXP1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ Jurkat cells were intracellularly stained with 0.4 ug Anti-Human FOXP1 (22051-1-AP) and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug x. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).