For Research Use Only

Cytokeratin 14 Polyclonal antibody

Catalog Number:22221-1-AP 2 Publications



Basic Information

Catalog Number: 22221-1-AP

Size:

AG17559

BC002690 GeneID (NCBI): 3861

GenBank Accession Number:

300 ug/ml **UNIPROT ID:** Source: P02533

Rabbit Full Name: Isotype: keratin 14 Calculated MW: Immunogen Catalog Number:

> 472 aa, 52 kDa Observed MW:

52 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:8000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF-P 1:50-1:500

Applications

Tested Applications: WB, IHC, IF-P, IP, ELISA

Cited Applications:

WB. IF

Species Specificity: human, mouse, rat **Cited Species:**

human, mouse, monkey, hamster

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse skin tissue, rat skin tissue

IP: mouse skin tissue.

IHC: human skin tissue, human bowen disease

IF-P: mouse skin tissue,

Background Information

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. Keratin 14 is a type I cytokeratin. It is usually found as a heterotetramer with keratin 5. Keratins K14 and K5 have long been considered to be biochemical markers of the stratified squamous epithelia, including epidermis. This antibody is specifically against KRT14.

Notable Publications

Author	Pubmed ID	Journal	Application
S Vijayalingam	24874842	Cancer Gene Ther	WB
Ming Zhou	39330987	Invest Ophthalmol Vis Sci	IF

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 22221-1-AP (Cytokeratin 14 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human skin tissue slide using 22221-1-AP (Cytokeratin 14 antibody at dilution of 1:200 (under 10) lengt




Immunohistochemical analysis of paraffinembedded human skin tissue slide using 22221-1-AP (Cytokeratin 14 antibody at dilution of 1:200 (under 60) lens)




IP result of anti-Cytokeratin 14 (IP:22221-1-AP, 4ug; Detection:22221-1-AP 1:1000) with mouse skin tissue lysate 3200ug.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse skin tissue using Cytokeratin 14 antibody (22221-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse skin tissue using Cytokeratin 14 antibody (22221-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human bowen disease slide using 22221-1-AP (Cytokeratin 14 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).