For Research Use Only

## Phospho-TDP43 (Ser409/410) Polyclonal antibody

Catalog Number:22309-1-AP 117 Publications



**Basic Information** 

Catalog Number: 22309-1-AP Source: Rabbit Isotype: GenBank Accession Number: NM\_007375 GeneID (NCBI): 23435 UNIPROT ID:

Q13148
Full Name:
TAR DNA bin

TAR DNA binding protein

Calculated MW: 43 kDa Observed MW: 40-50 kDa, 25-35 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:500-1:3000

**Applications** 

**Tested Applications:** 

WB, ELISA

Cited Applications: WB, IHC, IF, IP Species Specificity: human, mouse Cited Species:

human, mouse, rat, monkey

Positive Controls:

WB: Neuro-2a cells, Calyculin A treated HeLa cells

## **Background Information**

Transactivation response (TAR) DNA-binding protein of 43 kDa (also known as TARDBP or TDP-43) was first isolated as a transcriptional inactivator binding to the TAR DNA element of the HIV-1 virus. Neumann et al. (2006) found that a hyperphosphorylated, ubiquitinated, and cleaved form of TARDBP, known as pathologic TDP-43, is the major component of the tau-negative and ubiquitin-positive inclusions that characterize amyotrophic lateral sclerosis (ALS) and the most common pathological subtype of frontotemporal lobar degeneration (FTLD-U). Various forms of TDP-43 exist, including 18-35 kDa of cleaved C-terminal fragments, 45-50 kDa phospho-protein, 55 kDa glycosylated form, 75 kDa hyperphosphorylated form, and 90-300 kDa cross-linked form. (PMID: 17023659,19823856, 21666678, 22193176).22309-1-AP is a rabbit polyclonal antibody recognizing TDP-43 only when phosphorylated at 409/410. Immunohistochemical analyses using this antibody only stain the insoluble inclusions in pathologic tissues without normal diffuse nuclear staining.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Nikita Fernandes	32992901	Biomolecules	IF
Janice S W Ng	31529970	Biochemistry	IF
Ching-Chieh Chou	26130692	Hum Mol Genet	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**



Non-treated and Calyculin A treated HeLa cellls were subjected to SDS PAGE followed by western blot with 22309-1-AP (Phospho-TDP43 (Ser409/410) antibody) at dilution of 1:1000 incubated at room temperature for 1 hours.



Non-treated and Calyculin A treated Neuro-2a cellls were subjected to SDS PAGE followed by western blot with 22309-1-AP (Phospho-TDP43 (Ser409/410) antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.