For Research Use Only

LARG Polyclonal antibody

Catalog Number: 22441-1-AP

Featured Product

5 Publications



Purification Method:

WB 1:1000-1:4000 IHC 1:20-1:200

IF/ICC 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

Basic Information

Rho guanine nucleotide exchange

AG18211 Calculated MW: 1544 aa, 173 kDa Observed MW:

Observed 220 kDa

factor (GEF) 12

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IP Species Specificity:

Immunogen Catalog Number:

human Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: Jurkat cells, HeLa cells
IHC: human testis tissue,
IF/ICC: HepG2 cells, HeLa cells

Background Information

LARG, also known as ARHGEF12, is one of three regulator of G protein signalling (RGS) domain-containing RhoGEFs. LARG activates the ROCK pathway downstream of G α 12/13 signaling, leading to cytoskeletal reorganisation. LARG performs this activity either as a homodimer or as a heterodimer with Arhgef11. LARG has been implicated in many diseases, including cancer, heart disease, and in HIV viral replication. LARG is phosphorylated and dephosphorylated during mitosis and that Cdk1 is the mitotic kinase responsible for LARG phosphorylation. We got 220 kDa in western blotting, maybe due to its phosphorylation.

Notable Publications

Author	Pubmed ID	Journal	Application
Feng Zhang	33085791	Br J Pharmacol	WB
Hongyu Sun	36233059	Int J Mol Sci	WB
Zonghui Ding	32693062	Am J Pathol	WB,IHC

Storage

Storage:

Storage Ruffer

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.cor

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 22441-1-AP (LARG antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis slide using 22441-1-AP (LARG Antibody) at dilution of 1:50.



Immunohistochemical analysis of paraffinembedded human testis slide using 22441-1-AP (LARG Antibody) at dilution of 1:50.



Immunofluorescent analysis of HepG2 cells, using ARHGEF12 antibody 22441-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).