For Research Use Only

CNOT9 Polyclonal antibody

Catalog Number:22503-1-AP

Featured Product

15 Publications



Basic Information

Catalog Number: GenBank Accession Number: 22503-1-AP BC007102 GeneID (NCBI): Size: 367 µg/ml 9125 **UNIPROT ID:** Source:

Rabbit Q92600 Full Name: Isotype: RCD1 required for cell

Immunogen Catalog Number: Calculated MW:

AG18197

299 aa. 34 kDa Observed MW: 30-40 kDa

differentiation1 homolog (S. pombe)

Applications

Tested Applications: WB, IP, IF/ICC, IHC, ELISA **Cited Applications:**

WB, IP, CoIP Species Specificity: human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:500-1:2000 IF 1:20-1:200

Positive Controls:

WB: mouse thymus tissue, HepG2 cells, MCF-7 cells

IP: MCF-7 cells,

IHC: human stomach cancer tissue, rat colon tissue

IF: HepG2 cells,

Background Information

RQCD1, also known as CNOT9, is a highly conserved transcriptional cofactor predominantly expressed in actively diffierentiating tissues like testis and thymus. It is a component of the CCR4-NOT complex, the major effector complex in miRNA-mediated gene silencing. Overexpression of RQCD1 has been reported in serveral tumors, indicating that it may be a potential molecular target for cancer treatment.

Notable Publications

Author	Pubmed ID	Journal	Application
Alberto Carreño	35920669	Mol Cell Biol	WB,CoIP
Marta Mauri	27604873	Nucleic Acids Res	WB
Jessica Sheu-Gruttadauria	29576456	Cell	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




WB result of RQCD1 antibody (22503-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RQCD1 transfected HepG2 cells.




mouse thymus tissue were subjected to SDS PAGE followed by western blot with 22503-1-AP (RQCD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immun of luorescent analysis of HepG2 cells using 22503-1-AP (RQCD1 Antibody) at dilution of 1:50 and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-CNOT9 (IP:22503-1-AP, 4ug; Detection:22503-1-AP 1:1000) with MCF-7 cells lysate 1280ug.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 22503-1-AP (CNOT9 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 22503-1-AP (CNOT9 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).