

For Research Use Only

GAB2 Polyclonal antibody

Catalog Number: 22549-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number:

22549-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG18192

GenBank Accession Number:

BC131711

GeneID (NCBI):

9846

UNIPROT ID:

Q9UQC2

Full Name:

GRB2-associated binding protein 2

Calculated MW:

676 aa, 74 kDa

Observed MW:

74-90 kDa

Purification Method:

Antigen Affinity purified

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity:

human, mouse

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : K-562 cells,

IHC : mouse spleen tissue,

IF/ICC : K-562 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Lijun Mu	32901879	Oncol Rep	IHC
Jianxin Jiang	31737190	Am J Transl Res	WB
Xin Qian	33584322	Front Physiol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 22549-1-AP (GAB2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of GAB2 antibody (22549-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GAB2 transfected K-562 cells.



K-562 cells lysates were subjected to SDS PAGE followed by western blot with 22549-1-AP (GAB2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed K-562 cells using GAB2 antibody (22549-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).