For Research Use Only

## SART1 Polyclonal antibody

Catalog Number:22675-1-AP

Featured Product






Basic Information	Catalog Number: 22675-1-AP	GenBank Accession Number: BC001058		Purification Method: Antigen Affinity purified	
	Size:	GenelD (N	CBI):	Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:200-1:800 IF/ICC 1:400-1:1600	
	700 ug/ml	9092			
	Source:	UNIPROT I	D:		
	Rabbit	043290			
	Isotype: IgG	Full Name	: cell carcinoma antigen		
	Immunogen Catalog Number:	recognized	•		
	AG18478		Calculated MW: 800 aa, 90 kDa Observed MW: 110 kDa		
Applications	Tested Applications:		Positive Controls:		
	WB, IHC, IF/ICC, IP, ELISA		WB: HeLa cells, Jurkat cells		
	Cited Applications: WB, IF	B, IF		brain tissue,	
	Species Specificity:			testis tissue,	
	human, mouse		IF/ICC : HepG2 cells,		
	Cited Species: human, mouse				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	Squamous cell carcinoma antigen recognized by T cells (SART 1), also known as U4/U6.U5 tri-snRNP-associated 110 kDa protein, is a ubiquitously expressed protein that is involved in mRNA splicing and the regulation of cell proliferation. It has been identified as regulator of c-FLIP and drug-induced activation of caspase 8. The SART 1 gene encodes two proteins, the SART 1(800) protein is located in the nucleus of the majority of proliferating cells, while the SART1(259) protein is located in the cytosol of epithelial cancers. Genetic variation in the SART 1 gene may be associated with breast cancer development. This antibody recognizes the endogenous 800-amino acid SART 1, which migrates with an apparent molecular mass of 110 kDa in SDS-PAGE. (PMID: 22027693; 9449708; 19377877; 11350945)				
Notable Publications	Authory	Pulsus d1D	1	A surface to se	
	Author	Pubmed ID	Journal	Application	
	Ting Pan	33391530	Theranostics	IF,WB	
	Yalong Wang	38985674	Cell Rep	WB	
Storage	Storage: Store at -20°C. Stable for one ye Storage Buffer: PBS with 0.02% sodium azide at Aliquoting is unnecessary for -2	nd 50% glycerol j			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data


for 1.5 hours.




HeLa cells were subjected to SDS PAGE followed by western blot with 22675-1-AP (SART1 antibody) at dilution of 1:1000 incubated at room temperature




IP result of anti-SART1 (IP:22675-1-AP, 4ug; Detection:22675-1-AP 1:6000) with mouse brain tissue lysate 2100 ug.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 22675-1-AP (SART1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 22675-1-AP (SART1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepC2 cells using SART1 antibody (22675-1-AP) at dilution of 1:800 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red).