For Research Use Only

CCS Polyclonal antibody Catalog Number:22802-1-AP 9 Publications

Antibodies | ELISA kits | Proteins WWW.ptglab.com

Basic Information	Catalog Number: 22802-1-AP	GenBank Accession Number: BC105016		Purification Method: Antigen affinity purification			
	Size:	GeneID (NCBI)	:	Recommended Dilutions: WB 1:500-1:2000			
	900 µg/ml	9973					
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG18850	UNIPROT ID: IP 0.5-4.0 ug for 1.0-3.0 mg of 014618 protein lysate IHC 1:50-1:500 Full Name: IF/ICC 1:200-1:800 copper chaperone for superoxide dismutase Calculated MW: 274 aa, 29 kDa		IHC 1:50-1:500			
					Observed MW: 34 kDa	Observed MW: 34 kDa	
				Applications	Tested Applications: WB, IHC, IF/ICC, IP, ELISA	Positive Controls:	
		Cited Applications:			WB : HepG2 cells,		
WB, IHC							
Species Specificity: IHC : mouse brain tissue, human prostate cancer tissu human, mouse IF/ICC : HepG2 cells,							
Cited Species: human, mouse							
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0							
Background Information	Copper chaperone for superoxide dismutase (CCS) is a soluble factor that converts copper into copper-zinc superoxide dismutase (SOD1). CCS and SOD1 interact directly in vivo and in vitro, and this interaction is mediated by homologous domains in each protein. CCS not only interacts with wild-type SOD1 but also interacts with SOD1 containing common missense mutations, resulting in FALS.						
Notable Publications	Author	Pubmed ID	Journal	Application			
	Rashim Pal Singh	32416059	Cell Stem Cell	WB			
	Sarah E Antinone	28120938	Sci Rep	WB			
	Nadine Bakkar	33359139	Neurobiol Dis	WB			
Storage	Storage: Store at -20°C. Stable for one ye Storage Buffer: PBS with 0.02% sodium azide au Aliquoting is unnecessary for -2	nd 50% glycerol pH 7	·.3.				

 For technical support and original validation data for this product please contact:


 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 22802-1-AP (CCS antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-CCS (IP:22802-1-AP, 3ug; Detection:22802-1-AP 1:1000) with HepG2 cells lysate 6000ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 22802-1-AP (CCS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CCS antibody (22802-1-AP) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594phalloidin (red).