For Research Use Only

FOXC2 Polyclonal antibody

Catalog Number:23066-1-AP

Featured Product 16 Publications





Basic Information	Catalog Number: 23066-1-AP	GenBank Accession Number: BC113437	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	450 μ g/ml	2303	WB 1:500-1:1000 IF/ICC 1:50-1:500	
	Source: Rabbit	UNIPROT ID: Q99958	II / ICC 1.30-1.300	
	Isotype:	Full Name:		
	IgG Immunogen Catalog Number: AG19378	forkhead box C2 (MFH-1, mesenchyme forkhead 1)		
		Calculated MW: 501 aa, 54 kDa		
		Observed MW: 54-68 kDa		
Applications	Tested Applications:	Positive C	Positive Controls:	
	WB, IF/ICC, ELISA	WB: HCT	WB : HCT 116 cells, rat spleen tissue, MCF-7 cells	
	Cited Applications: WB, IHC, IF	IF/ICC : A	IF/ICC : A375 cells,	
	Species Specificity: human, mouse, rat			
	Cited Species: human, mouse, rat			
Background Information	Forkhead box protein C2 (FOXC2) also known as forkhead-related protein FKHL14 (FKHL14), transcription factor 14, or mesenchyme fork head protein 1 (MFH1) is a protein that in humans is encoded by the FOXC2 gene. FOXC a member of the fork head box (FOX) family of transcription factors. FOX transcription factors are expressed duri development and are associated with a number of cellular and developmental differentiation processes. FOXC2 required during early development of the kidneys, including differentiation of podocytes and maturation of the glomerular basement membrane. It is also involved in the early development of the heart. FOXC2 is also involved in the early development of the heart.			
	development and are associated required during early development	nt of the kidneys, including differentia	tion of podocytes and maturation of the	
	development and are associated or required during early development glomerular basement membrane. in cancer metastases. In particula mesenchymal transition (EMT) and	nt of the kidneys, including differentia It is also involved in the early develo r, expression of FOXC2 is induced whe d become mesenchymal looking cell OXC2 protein, but the phosphorylated	tion of podocytes and maturation of the poment of the heart. FOXC2 is also involven epithelial cells undergo an epithelial-	
	development and are associated or required during early development glomerular basement membrane. in cancer metastases. In particula mesenchymal transition (EMT) an antibody recognizes the 56 kDa Fi 62kDa band was also detected in	nt of the kidneys, including differentia It is also involved in the early develo r, expression of FOXC2 is induced whe d become mesenchymal looking cell OXC2 protein, but the phosphorylated	tion of podocytes and maturation of the opment of the heart. FOXC2 is also involve en epithelial cells undergo an epithelial- 5. (PMID: 8674414 9169153 19935708) Thi	
	development and are associated or required during early development glomerular basement membrane. in cancer metastases. In particula mesenchymal transition (EMT) an antibody recognizes the 56 kDa Fe 62kDa band was also detected in Author	nt of the kidneys, including differentia It is also involved in the early develo r, expression of FOXC2 is induced whe d become mesenchymal looking cells OXC2 protein, but the phosphorylated the study(PMID: 19540201).	ntion of podocytes and maturation of the opment of the heart. FOXC2 is also involven epithelial cells undergo an epithelial- 5. (PMID: 8674414 9169153 19935708) Thi FOXC2 may be a 56-65 kDa protein. The	
	development and are associated or required during early development glomerular basement membrane. in cancer metastases. In particula mesenchymal transition (EMT) an antibody recognizes the 56 kDa Ft 62kDa band was also detected in Author Jianwen Liao	nt of the kidneys, including differentia It is also involved in the early develor r, expression of FOXC2 is induced whe d become mesenchymal looking cells OXC2 protein, but the phosphorylated the study(PMID: 19540201).	tion of podocytes and maturation of the opment of the heart. FOXC2 is also involve en epithelial cells undergo an epithelial- 5. (PMID: 8674414 9169153 19935708) Thi FOXC2 may be a 56-65 kDa protein. The Application	
Notable Publications	development and are associated a required during early development glomerular basement membrane. in cancer metastases. In particula mesenchymal transition (EMT) an antibody recognizes the 56 kDa Fe 62kDa band was also detected in Author I Jianwen Liao Eei-Xiang Lin	nt of the kidneys, including differentia It is also involved in the early develor, expression of FOXC2 is induced whe d become mesenchymal looking cells DXC2 protein, but the phosphorylated the study(PMID: 19540201). Pubmed ID Journal 36106596 J Neurotrauma	tion of podocytes and maturation of the opment of the heart. FOXC2 is also involve en epithelial cells undergo an epithelial- 5. (PMID: 8674414 9169153 19935708) Thi FOXC2 may be a 56-65 kDa protein. The Application WB	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 23066-1-AP (FOXC2 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours. Immunofluorescent analysis of (4% PFA) fixed A375 cells using 23066-1-AP (FOXC2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).