## For Research Use Only

# Alpha Actin Polyclonal antibody

Catalog Number:23082-1-AP 1 Publications



**Basic Information** 

Catalog Number: 23082-1-AP

GenBank Accession Number: BC009978

Antigen affinity purification

Size: 500 ug/ml GeneID (NCBI): 70 Recommended Dilutions: WB 1:500-1:2000

Source: Rabbit UNIPROT ID: P68032 IHC 1:200-1:1000

**Purification Method:** 

Isotype: Full Name:
IgG actin, alpha, cardiac muscle 1

Immunogen Catalog Number: Calculated MW:

AG19482 377 aa, 42 kDa

Observed MW: 42-45 kDa

**Applications** 

Tested Applications:

WB, IHC, FC (Intra), ELISA

Cited Applications:

WB

Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse heart tissue,

IHC: human normal colon, human colon tissue, human heart tissue, human skeletal muscle tissue, human small intestine tissue, mouse skeletal muscle tissue, mouse heart tissue, mouse colon tissue

# **Background Information**

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Hideshi Okada	25466765	Circ Heart Fail	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**



mouse heart tissue were subjected to SDS PAGE followed by western blot with 23082-1-AP (alpha Actin antibody at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human heart using 23082-1-AP (alpha Actin antibody) at dilution of 1:100 (under 10x lens)



Immunohistochemical analysis of paraffinembedded human heart using 23082-1-AP (alpha Actin antibody) at dilution of 1:100 (under 40x lens).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 23082-1-AP (Alpha Actin antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 C2C12 cells were intracellularly stained with 0.4 ug Alpha Actin Polyclonal antibody (23082-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).