### For Research Use Only

# MYD88 Polyclonal antibody

Catalog Number:23230-1-AP

Featured Product

212 Publications



**Basic Information** 

Catalog Number:

23230-1-AP

Concentration:

600 µg/ml

Source:

Rabbit

GenBank Accession Number:

BC013589

GeneID (NCBI):

4615

UNIPROT ID:

Rabbit

Q99836

Isotype:

Full Name:

myeloid differentiation primary

Immunogen Catalog Number: response gene (88)
AG19770 Calculated MW:
33 kDa

Observed MW: 33 kDa

**Applications** 

Tested Applications: WB, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP

Species Specificity: human Cited Species:

human, pig, canine, monkey, chicken, bovine, sheep

**Positive Controls:** 

WB: Jurkat cells, A549 cells, HepG2 cells, MCF-7 cells,

**Purification Method:** 

WB 1:5000-1:50000

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

K-562 cells, Raji cells

IP: K-562 cells, A549 cells

## **Background Information**

MYD88 is a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. MYD88 is the key adaptor protein for the interleukin-1 receptor (IL-1R) and most Toll-like receptors (TLRs), which are essential for innate immunity and pathogen-associated molecular pattern recognition. Mutations in the MyD88 gene lead to the development of cancer in humans and mice suggesting that MyD88 also plays a cell autonomous role in tissue homeostasis.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Chang Liu	36230117	Foods	WB
Yangming Zhang	36191372	Biochem Biophys Res Commun	WB
Zhaohai Wen	36238295	Front Immunol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffe


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


#### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 23230-1-AP (MYD88 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




WB result of MYD88 antibody (23230-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MYD88 transfected HepG2 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 23230-1-AP (MYD88 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-MYD88 (IP:23230-1-AP, 4ug; Detection:23230-1-AP 1:10000) with K-562 cells lysate 1360 ug.



IP result of anti-MYD88 (IP:23230-1-AP, 4ug; Detection:23230-1-AP 1:10000) with A549 cells lysate 640 ug.