For Research Use Only

KCNN4 Polyclonal antibody

Catalog Number:23271-1-AP

Featured Product

13 Publications



Basic Information

Catalog Number: 23271-1-AP Source: Rabbit

Isotype: IgG

Immunogen Catalog Number:

AG19774

GenBank Accession Number:

BC015337 GeneID (NCBI): 3783 UNIPROT ID: 015554

potassium intermediate/small conductance calcium-activated channel, subfamily N, member 4

Calculated MW: 427 aa, 48 kDa Observed MW: 48 kDa

Full Name:

Purification Method: Antigen Affinity purified Recommended Dilutions: WB: 1:1000-1:8000

IF/ICC: 1:200-1:800

Applications

Tested Applications: WB, IF/ICC, ELISA Cited Applications: WB, IHC, IF, IP, CoIP

Species Specificity: human

Cited Species: human, mouse, canine **Positive Controls:**

WB: A431 cells, HepG2 cells, HEK-293 cells

IF/ICC: HEK-293 cells,

Background Information

The intermediate conductance calcium-activated potassium channel KCNN4, also known as KCa3.1, IK1, or SK4, is a member of the potassium channel KCNN family. KCNN4 is expressed in T cells, mast cells, macrophages, erythrocytes, vascular smooth muscle cells, airway smooth muscle cells, intestinal and airway epithelial cells (PMID: 19681865; 21493782). Functionally, KCNN4 plays a key role in calcium-dependent cell functions, such as proliferation, activation, and migration, in a broad range of cell types (PMID: 21493782).

Notable Publications

Author	Pubmed ID	Journal	Application
Shuheng Jiang	29050937	Biochem Biophys Res Commun	IHC
Zilin Qiao	36423196	Viruses	WB
Yankang Cui	35689211	Cancer Cell Int	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



WB result of KCNN4 antibody (23271-1-AP, 1:4000) with si-control and si-KCNN4 transfected HEK293 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 23271-1-AP (KCNN4 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using KCNN4 antibody (23271-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).