For Research Use Only

SGK1 Polyclonal antibody Catalog Number:23394-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: 23394-1-AP	GenBank Accession Number: BC001263	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	1000 µg/ml	6446	IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Source:	UNIPROT ID:	protein lysate	
	Rabbit	000141	IHC 1:20-1:200	
	Isotype:	Full Name:	IF/ICC 1:50-1:500	
	lgG	serum/glucocorticoid regulated	cocorticoid regulated	
	Immunogen Catalog Number: AG17888	kinase 1		
		Calculated MW: 431 aa, 49 kDa		
Applications	Tested Applications:	Positive	Positive Controls:	
	IHC, IF/ICC, IP, ELISA	IP : HEK-2	93 cells,	
	Cited Applications: IHC	IHC : hun	aan pancreas tissue, human breast cancer	
	Species Specificity:	tissue		
	human, mouse	IF/ICC : I	IEK-293 cells,	
	Cited Species: human, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	Serum- and glucocorticoid-induced kinase 1 (SGK1), also named as SGK, is a S/T protein kinase that belongs to the AGC (cAMP-dependent, cGMP-dependent, and protein kinase C) kinase family. It is expressed in many cell types a participates in numerous cellular processes and it is ubiquitinated and degraded at the ER membrane(PMID: 16847254). The N-terminal motif of SGK1 is critical for its ubiquitination and degradation(PMID:16817852). SGK a Akt are likely to phosphorylate related substrates, as they share a similar consensus phosphorylation site (RXRXXS/T)(PMID:11154281). It has 5 isoforms produced by alternative promoter usage and alternative splicing.			
Background Informatior	participates in numerous cellular 16847254). The N-terminal motif Akt are likely to phosphorylate re	processes and it is ubiquitinated and of SGK1 is critical for its ubiquitinatic lated substrates, as they share a simi	family. It is expressed in many cell types degraded at the ER membrane(PMID: in and degradation(PMID:16817852). SGK lar consensus phosphorylation site	
	participates in numerous cellular 16847254). The N-terminal motif Akt are likely to phosphorylate re (RXRXXS/T)(PMID:11154281). It ha	processes and it is ubiquitinated and of SGK1 is critical for its ubiquitinatic lated substrates, as they share a simi	family. It is expressed in many cell types degraded at the ER membrane(PMID: in and degradation(PMID:16817852). SGK lar consensus phosphorylation site promoter usage and alternative splicing.	
	AGC (CAMP-dependent, COMP-dep participates in numerous cellular 16847254). The N-terminal motif Akt are likely to phosphorylate re (RXRWS/T)(PMID:11154281). It ha	processes and it is ubiquitinated and of SGK1 is critical for its ubiquitinatio lated substrates, as they share a simi as 5 isoforms produced by alternative Pubmed ID Journal	family. It is expressed in many cell types degraded at the ER membrane(PMID: in and degradation(PMID:16817852). SGK lar consensus phosphorylation site	
	Add (CAMP-dependent, Camp-dep participates in numerous cellular 16847254). The N-terminal motif Akt are likely to phosphorylate re (RXRXXS/T)(PMID:11154281). It ha Author Huaxin Zhou	processes and it is ubiquitinated and of SGK1 is critical for its ubiquitinatio lated substrates, as they share a simi is 5 isoforms produced by alternative Pubmed ID Journal	family. It is expressed in many cell types degraded at the ER membrane(PMID: in and degradation(PMID:16817852). SGK lar consensus phosphorylation site promoter usage and alternative splicing.	
Background Informatior Notable Publications	Add (CAMP-dependent, ComP-dep participates in numerous cellular 16847254). The N-terminal motif Akt are likely to phosphorylate re (RXRXXS/T)(PMID:11154281). It ha Author Huaxin Zhou Junling Ren	processes and it is ubiquitinated and of SGK1 is critical for its ubiquitinatic lated substrates, as they share a simi is 5 isoforms produced by alternative Pubmed ID Journal 25993992 FASEB J	family. It is expressed in many cell types degraded at the ER membrane(PMID: in and degradation(PMID:16817852). SGK lar consensus phosphorylation site promoter usage and alternative splicing.	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human pancreas slide using 23394-1-AP (SGK1 Antibody) at dilution of 1:50.



Immunohistochemical analysis of paraffinembedded human pancreas slide using 23394-1-AP (SGK1 Antibody) at dilution of 1:50.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using SGK1 antibody (23394-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-SGK1 (IP:23394-1-AP, 4ug; Detection:23394-1-AP 1:600) with HEK-293 cells lysate 1480ug.