For Research Use Only

P15RS Polyclonal antibody

Catalog Number:23652-1-AP

Featured Product

4 Publications



Basic Information

Catalog Number: 23652-1-AP Size: 800 µ g/ml Source: Rabbit

gG

Immunogen Catalog Number:

AG20409

Isotype:

GenBank Accession Number:

BC010136
GeneID (NCBI):
55197
UNIPROT ID:
Q96P16
Full Name:

regulation of nuclear pre-mRNA domain containing 1A

Calculated MW:

312 aa, 36 kDa Observed MW: 36 kDa Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:6000 IF/ICC 1:50-1:500

Applications

Tested Applications: WB, IF/ICC, ELISA Cited Applications: WB, IHC, IP, IF Species Specificity: human, mouse Cited Species: human, mouse **Positive Controls:**

WB: A431 cells, mouse liver tissue, mouse skeletal muscle tissue, LO2 cells, Jurkat cells, HEK-293 cells

IF/ICC: HeLa cells, HepG2 cells

Background Information

RPRD1A is upregulated in cells overexpressing cyclin-dependent kinase inhibitor p15(INK4b) and may have a role in cell cycle regulation. It contains a RAR domain that is involved in regulation of nuclear pre-mRNA, which suggests that P15RS may act as a nuclear regulation protein [PMID:12470661]. It also interacts with phosphorylated C-terminal heptapeptide repeat domain (CTD) of the largest RNA polymerase II subunit POLR2A, and participates in dephosphorylation of the CTD. Besides, RPRD1A regulates cell cycle genes and attenuates WNT signaling [PMID: 20739273], and acts as a negative regulator of cyclin-D1(CCND1) and cyclin-E (CCNE1) in the cell cycle [PMID: 22231121].

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaoman Chen	30207732	Mol Pharm	WB
Simona Cugusi	36121223	Mol Cell Biol	WB
Xiaofan Feng	34921137	Cell Death Dis	IP,WB,IHC,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 23652-1-AP (P15RS antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 23652-1-AP (P15RS antibody), at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).