For Research Use Only

AIDA Polyclonal antibody

Catalog Number:23724-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number:

23724-1-AP

BC015535

Size:

GeneID (NCBI):

800 µg/ml

64853

Source:

UNIPROT ID:

Rabbit

Q96BJ3

Isotype:

GeneBank Accession Number:

BC015535

GeneID (NCBI):

GeneID (NCBI):

GeneID (NCBI):

Full Name:

IgG axin interactor, dorsalization

Immunogen Catalog Number: associated
AG20588 Calculated MW:
306 aa, 35 kDa

Observed MW: 35-40 kDa Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000

Purification Method:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

Applications

Tested Applications: IHC, IP, WB,ELISA Cited Applications: WB, IP, IF Species Specificity: human, mouse

Cited Species:
mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293 cells,
IP: HeLa cells,

IHC: mouse skeletal muscle tissue, human heart

tissue

Background Information

AIDA, also named as C1orf80, acts as a ventralizing factor during embryogenesis. It inhibits axin-mediated JNK activation by binding axin and disrupting axin homodimerization. It's highest expression in the heart and skeletal muscle. AIDA has three isoforms with MW 35 kDa, 16 kDa and 26 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Meng Shi	33664495	Nat Cell Biol	IP
Hui Luo	29617643	Cell Metab	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 23724-1-AP (AIDA antibody) at dilution of 1:100 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 23724-1-AP (AIDA antibody) at dilution of 1:100 (under 0x lens).



HEK-293 cells were subjected to SDS PAGE followed by western blot with 23724-1-AP (AIDA Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-AIDA (IP:23724-1-AP, 3ug; Detection:23724-1-AP 1:600) with HeLa cells lysate 4000ug.