For Research Use Only

SNRNP200 Polyclonal antibody

Catalog Number: 23875-1-AP

3 Publications



Basic Information

Catalog Number: 23875-1-AP Size: 400 μg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG20748

GenBank Accession Number:

BC001417 GeneID (NCBI): 23020 **UNIPROT ID:** 075643 Full Name:

small nuclear ribonucleoprotein 200kDa (U5)

Calculated MW: 494 aa, 57 kDa Observed MW: 200-245 kDa

Purification Method: Antigen Affinity purified Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

Applications

Tested Applications: FC, IP, WB, ELISA Cited Applications:

Species Specificity: human, mouse Cited Species: human

Positive Controls:

WB: HeLa cells, NIH/3T3 cells, HEK-293 cells, Jurkat

cells

IP: HeLa cells,

Background Information

SNRNP200 (small nuclear ribonucleoprotein 200kDa (U5)), also known as HELIC2, ASCC3L1 or BRR2, is a 2,136 amino $acid\ protein\ that\ localizes\ to\ the\ nucleus\ and\ contains\ two\ SEC63\ domains, two\ helicase\ C-terminal\ domains\ and$ two helicase ATP-binding domains. Existing as multiple alternatively spliced isoforms, HELIC2 functions as an RNA helicase that is thought to promote specific RNA-RNA conformational changes which are important in the second step of RNA splicing. The gene encoding HELIC2 maps to human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome. Harlequin icthyosis, a rare and morbid skin deformity, is associated with mutations in the chromosome 2-localized ABCA12 gene, while the lipid metabolic disorder sitosterolemia is associated with defects in the ABCG5 and ABCG8 genes, hich also map to chromosome 2. The observed molecular weight of SNRNP200 is 200-245kd.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhe Chen	28878014	J Biol Chem	IP
Shuai Hou	31216022	Nucleic Acids Res	WB
Bangming Jin	37395406	Nucleic Acids Res	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 23875-1-AP (SNRNP200 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-SNRNP200 (IP:23875-1-AP, 4ug; Detection:23875-1-AP 1:1000) with HeLa cells lysate 4000ug.



Various lysates were subjected to SDS PAGE followed by western blot with 23875-1-AP (SNRNP200 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



1X10^6 A431 cells were intracellularly stained with 0.4 ug Anti-Human SNRNP200 (23875-1-AP) and Coralite®647-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).