## For Research Use Only

# SOX7 Polyclonal antibody

Catalog Number: 23925-1-AP 5 Publications



**Basic Information** 

Catalog Number:
23925-1-AP
BC071947
Size:
GeneID (NCBI):
750 ug/ml
83595
Source:
UNIPROT ID:
Rabbit
Q9BT81
Isotype:
Full Name:

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500 IF/ICC 1:200-1:800

Isotype:

SRY (sex determining region Y)-box 7

Immunogen Catalog Number:Calculated MW:AG10323388 aa, 42 kDaObserved MW:

42-45 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IF, IP

Species Specificity: human, mouse Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: COLO 320 cells, mouse heart tissue, PC-3 cells IHC: human prostate cancer tissue, human colon tissue

IF/ICC: U2OS cells,

# **Background Information**

SOX7, also named as Transcription factor SOX-7, is a 388 amino acid protein, which is widely expressed in adult and fetal tissues and tends to be down-regulated in prostate adenocarcinomas and colorectal tumors due to promoter hypermethylation. SOX7 binds to and activates the CDH5 promoter, hence plays a role in the transcriptional regulation of genes expressed in the hemogenic endothelium and blocks further differentiation into blood precursors. SOX7 may be required for the survival of both hematopoietic and endothelial precursors during specification.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Baolei Li	35260939	Mol Genet Genomics	WB,IP
Qiao-Yang Sun	31332289	Oncogene	WB
Nanchao Hong	33846290	Cell Death Dis	WB

# Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 23925-1-AP (SOX7 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 23925-1-AP (SOX7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 23925-1-AP (SOX7 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using SOX7 antibody (23925-1-AP) at dilution of 1:400 and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).