For Research Use Only

MAP4K1 Polyclonal antibody

Catalog Number:23950-1-AP

3 Publications



Basic Information

Catalog Number:
23950-1-AP
BC152994
Size:
GeneID (NCBI):
500 µ g/ml
11184
Source:
Rabbit
Q92918
Isotype:
GeneID (NCBI):
Full Name:

mitogen-activated protein kinase

90 kDa

Calculated MW:

Immunogen Catalog Number: kinase kinase kinase 1

833 aa, 91 kDa Observed MW:

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications:

AG20221

Species Specificity:

human Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: Jurkat cells, Raji cells

IP: Jurkat cells,

IHC: human tonsillitis tissue, human liver tissue

Purification Method:

WB 1:1000-1:8000

protein lysate

IHC 1:50-1:500

Antigen Affinity purified

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Background Information

MAP4K1(Mitogen-activated protein kinase kinase kinase kinase 1) is also named as MEKKK 1, HPK1 and belongs to the protein kinase superfamily. It may function as a novel tumor suppressor and its loss plays a critical role in pancreatic cancer. MAP4K1 regulates stress responses, proliferation, and apoptosis in hematopoietic cells. This protein has 2 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Xunping Zhao	34530056	Cell Signal	WB
Li Zhou	37495569	Nat Commun	WB
Lin Yang	37106265	Clin Exp Med	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 23950-1-AP (MAP4K1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 23950-1-AP (MAP4K1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-MAP4K1 (IP:23950-1-AP, 4ug; Detection:23950-1-AP 1:1000) with Jurkat cells lysate 3200ug.



Various lysates were subjected to SDS PAGE followed by western blot with 23950-1-AP (MAP4K1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 23950-1-AP (MAP4K1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 23950-1-AP (MAP4K1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).