For Research Use Only

RCOR2 Polyclonal antibody Catalog Number:23969-1-AP Featured Product







	Catalog Number: 23969-1-AP	GenBank Accession Numbe BC023587	r: Purification Method: Antigen Affinity purified	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	300 µg/ml Source: Rabbit	283248 UNIPROT ID: 08IZ40	WB 1:500-1:2000	
			IHC 1:50-1:500	
	Isotype:	Full Name:		
	lgG	REST corepressor 2		
	Immunogen Catalog Number: AG21112	Calculated MW: 523 aa, 58 kDa		
		Observed MW: 58 kDa		
Applications	Tested Applications:	Pos	Positive Controls:	
	WB, IHC, ELISA Cited Applications:	WB	WB : Jurkat cells, HEK-293 cells	
	WB, ColP	IHC	IHC : mouse embryo tissue,	
	Species Specificity: human, mouse			
	Cited Species: human, mouse			
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0	atively, antigen		
	In mammals, the CoREST (corepressor for element-1-silencing transcription factor) complex is a chromatin- modifying structure that, through interactions with NRSF (neuron restrictive silencer factor), regulates neuronal ger expression and neuronal cell fate. RCOR2 (REST corepressor 2) and RCOR3 (REST corepressor 3) are nuclear proteins that each contain one ELM2 domain and two SANT domains. RCOR2 and RCOR3, both members of the CoREST family, are thought to function as components of the CoREST complex, possibly playing a role in the transcriptional repression of neuronal genes. Additionally, RCOR2 and RCOR3, in conjunction with CoREST, can form immunocomplexes with a variety of histone-modifying genes, including G9a and HDAC1. Via these protein complexes, RCOR2 and RCOR3 can further regulate transcription by controlling the methylation and demethylatior of target genes during early development.			
Background Information	modifying structure that, through expression and neuronal cell fate that each contain one ELM2 doma family, are thought to function as repression of neuronal genes. Add immunocomplexes with a variety complexes, RCOR2 and RCOR3 ca	RCOR2 (REST corepressor 2) an in and two SANT domains. RCO components of the CoREST com litionally, RCOR2 and RCOR3, ir of histone-modifying genes, in n further regulate transcription	restrictive silencer factor), regulates neuronal ge d RCOR3 (REST corepressor 3) are nuclear protei R2 and RCOR3, both members of the CoREST plex, possibly playing a role in the transcription conjunction with CoREST, can form cluding G9a and HDAC1. Via these protein	
	modifying structure that, through expression and neuronal cell fate that each contain one ELM2 doma family, are thought to function as repression of neuronal genes. Add immunocomplexes with a variety complexes, RCOR2 and RCOR3 ca of target genes during early deve	RCOR2 (REST corepressor 2) an in and two SANT domains. RCO components of the CoREST com litionally, RCOR2 and RCOR3, ir of histone-modifying genes, in n further regulate transcription l opment.	restrictive silencer factor), regulates neuronal ge d RCOR3 (REST corepressor 3) are nuclear protein 22 and RCOR3, both members of the CoREST plex, possibly playing a role in the transcription a conjunction with CoREST, can form cluding G9a and HDAC1. Via these protein by controlling the methylation and demethylation	
	modifying structure that, through expression and neuronal cell fate that each contain one ELM2 doma family, are thought to function as repression of neuronal genes. Add immunocomplexes with a variety complexes, RCOR2 and RCOR3 ca of target genes during early deve	RCOR2 (REST corepressor 2) an in and two SANT domains. RCO components of the CoREST com litionally, RCOR2 and RCOR3, ir of histone-modifying genes, in n further regulate transcription I opment.	restrictive silencer factor), regulates neuronal ge d RCOR3 (REST corepressor 3) are nuclear protei 22 and RCOR3, both members of the CoREST plex, possibly playing a role in the transcription conjunction with CoREST, can form cluding G9a and HDAC1. Via these protein by controlling the methylation and demethylation	
	modifying structure that, through expression and neuronal cell fate that each contain one ELM2 doma family, are thought to function as repression of neuronal genes. Add immunocomplexes with a variety complexes, RCOR2 and RCOR3 ca of target genes during early deve Author F Donglim Esther Park	RCOR2 (REST corepressor 2) an in and two SANT domains. RCO components of the CoREST com litionally, RCOR2 and RCOR3, ir of histone-modifying genes, in n further regulate transcription l copment.	restrictive silencer factor), regulates neuronal ge d RCOR3 (REST corepressor 3) are nuclear protei 22 and RCOR3, both members of the CoREST plex, possibly playing a role in the transcription is conjunction with CoREST, can form cluding G9a and HDAC1. Via these protein by controlling the methylation and demethylation Application WB	
Background Information	modifying structure that, through expression and neuronal cell fate that each contain one ELM2 doma family, are thought to function as repression of neuronal genes. Add immunocomplexes with a variety complexes, RCOR2 and RCOR3 ca of target genes during early deve Author F Donglim Esther Park Sandhya Malla	RCOR2 (REST corepressor 2) and in and two SANT domains. RCOI components of the CoREST com litionally, RCOR2 and RCOR3, ir of histone-modifying genes, in n further regulate transcription loopment. Pubmed ID Journal 2284543 Nat Cell Biol 9237615 Nat Communication	restrictive silencer factor), regulates neuronal ge d RCOR3 (REST corepressor 3) are nuclear protein 22 and RCOR3, both members of the CoREST plex, possibly playing a role in the transcription conjunction with CoREST, can form cluding G9a and HDAC1. Via these protein by controlling the methylation and demethylation Application WB n WB,CoIP	
	modifying structure that, through expression and neuronal cell fate that each contain one ELM2 doma family, are thought to function as repression of neuronal genes. Add immunocomplexes with a variety complexes, RCOR2 and RCOR3 ca of target genes during early deve Author F Donglim Esther Park Sandhya Malla	RCOR2 (REST corepressor 2) an in and two SANT domains. RCO components of the CoREST com litionally, RCOR2 and RCOR3, ir of histone-modifying genes, in n further regulate transcription l copment.	restrictive silencer factor), regulates neuronal ge d RCOR3 (REST corepressor 3) are nuclear protei 22 and RCOR3, both members of the CoREST plex, possibly playing a role in the transcription conjunction with CoREST, can form cluding G9a and HDAC1. Via these protein by controlling the methylation and demethylation by controlling the methylation with CoREST WB n WB,CoIP	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Jurkat cells were subjected to SDS PAGE followed by western blot with 23969-1-AP (RCOR2 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse embryo tissue slide using 23969-1-AP (RCOR2 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).