## For Research Use Only

# Sodium iodide symporter Polyclonal antibody



**Purification Method:** 

WB 1:500-1:1000

protein lysate

IHC 1:20-1:200

Positive Controls:

IP: mouse testis tissue,

7901 cells

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

WB: mouse testis tissue, mouse stomach tissue, SGC-

IHC: human thyroid cancer tissue, human ovary tissue

Catalog Number: 24324-1-AP

**40 Publications** 

**Basic Information** 

Catalog Number:

24324-1-AP

BC105047

Size:

GeneID (NCBI):

1200 µg/ml

6528

Source:

UNIPROT ID:

Rabbit

Q92911

Isotype:

GenBank Accession Number:

BC105047

GeneID (NCBI):

GeneID (NCBI):

Full Name:

gG solute carrier family 5 (sodium iodide symporter), member 5

Immunogen Catalog Number: symporter), member: AG19504 Calculated MW:

643 aa, 69 kDa Observed MW: 50-55 kDa, 75-100 kDa

**Applications** 

Tested Applications: FC, IHC, IP, WB, ELISA Cited Applications: WB, IP, IF, FC, IHC, ChIP Species Specificity: human, mouse, rat Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

# **Background Information**

The sodium iodide symporter (Na+/I - symporter, NIS), encoded by SLC5A5, is an integral plasma membrane glycoprotein that plays an important role in iodide uptake by thyroid cells. Expression of sodium iodide symporter has also been found in extra-thyroidal tissues, including gastric mucosa, lactating mammary gland and salivary glands. Increased expression of sodium iodide symporter has been found in thyroid tissue from patients with Graves' disease as well as papillary thyroid carcinomas. In addition, sodium iodide symporter was found to express in majority of breast cancer tissue but not in normal tissue. Sodium iodide symporter can be a promising diagnostic and therapeutic tool for thyroid cancer and breast cancer. This antibody recognizes the mature approximately 75-100 kDa protein and a partially glycosylated 50-55 kDa protein. (PMID: 12588808, 9525971)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jun Wang	32940055	Hum Gene Ther	IF
Martin L Read	34520744	Cell Chem Biol	WB
Jianlu Song	30323976	Am J Cancer Res	WB

### Storage

Storage


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




mouse testis tissue were subjected to SDS PAGE followed by western blot with 24324-1-AP (Sodium iodide symporter antibody at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 24324-1-AP (Sodium iodide symporter antibody at dilution of 1:50 (under 40x lens).



1X10^6 MCF-7 cells were stained with 0.2ug Sodium iodide symporter antibody (24324-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:100.



IP result of anti-Sodium iodide symporter (IP:24324-1-AP, 4ug; Detection:24324-1-AP 1:500) with mouse testis tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 24324-1-AP (Sodium iodide symporter antibody at dilution of 1:50 (under 10x lens).