## For Research Use Only

# TOP2A Polyclonal antibody

Catalog Number: 24641-1-AP

Featured Product

15 Publications



**Basic Information** 

Catalog Number: 24641-1-AP

Size:

1000 µg/ml

GenBank Accession Number: BC140791

GeneID (NCBI): 7153

Source: UNIPROT ID:

Rabbit P11388
Isotype: Full Name:

lgG topoisomerase (DNA) II alpha 170kDa IF/ICC 1:50-1:500

Immunogen Catalog Number: Calculated MW:

AG19551 1531 aa, 174 kDa

Observed MW: 174 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity:

human, mouse
Cited Species:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: HEK-293T cells, HeLa cells, HepG2 cells, MDA-MB-

**Purification Method:** 

WB 1:2000-1:16000

protein lysate

IHC 1:50-1:500

Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

231 cells

IP: HeLa cells,

IHC: human pancreas cancer tissue, human breast

IF/ICC : A431 cells,

cancer tissue

## **Background Information**

TOP2A, also named as TOP2, belongs to the type II topoisomerase family. Topoisomerase II makes double-strand breaks. TOP2 control the topological states of DNA by transient breakage and subsequent rejoining of DNA strands. TOP2A may play a role in regulating the period length of ARNTL/BMAL1 transcriptional oscillation. DNA topoisomerase IIa (170 kDa, TOP2a/170) is essential in proliferating cells by resolving DNA topological entanglements during chromosome condensation, replication, and segregation.

#### Notable Publications

Author	Pubmed ID	Journal	Application
Jun-Song Liu	25333415	Oncol Rep	
Jingjun Wu	26612655	Chem Biol Interact	WB
Li Wang	33236014	bioRxiv	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


#### **Selected Validation Data**




WB result of TOP2A antibody (24641-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TOP2A transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 24641-1-AP (TOP2A antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP result of anti-TOP2A (IP:24641-1-AP, 4ug; Detection:24641-1-AP 1:1000) with HeLa cells lysate 2000ug.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using TOP2A antibody (24641-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 24641-1-AP (TOP2A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).