## For Research Use Only

# NRBF2 Polyclonal antibody

Catalog Number: 24858-1-AP

Featured Product

5 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 24858-1-AP BC001345 GeneID (NCBI): Size: 1500 ug/ml 29982 **UNIPROT ID:** Source: Rabbit Q96F24

Isotype: nuclear receptor binding factor 2

Calculated MW: Immunogen Catalog Number:

AG21403 32 kDa

> Observed MW: 35-37 kDa

Full Name:

**Purification Method:** Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, IF, CoIP Species Specificity: human, mouse **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HeLa cells

IP: HEK-293 cells, IHC: human liver tissue, IF/ICC: HepG2 cells,

# **Background Information**

NRBF2 also termed as Nuclear receptor-binding factor 2or COPR is a 287 amino acid protein, which Interacts with PPARA, PPARD and PPARG. NRBF2 as transcription activator may modulate transcriptional activation by target nuclear receptors. NRBF2 is expressed in keratinocytes, liver and placenta. NRBF2, which exists as two isoforms due to alternative splicing, is localized to both the nucleus and the cytoplasm.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Hanhai Zeng	34530854	J Neuroinflammation	IHC,IF
Weitao Shen	35712081	iScience	WB,IHC,IF,CoIP
Xiaosen Ouyang	32350405	Lab Invest	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 24858-1-AP (NRBF2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 24858-1-AP (NRBF2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using NRBF2 antibody (24858-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP result of anti-NRBF2 (IP:24858-1-AP, 4ug; Detection:24858-1-AP 1:2000) with HEK-293 cells lysate 2640ug.