For Research Use Only

POGK Polyclonal antibody

Catalog Number:24888-1-AP




Catalog Number: 24888-1-AP	GenBank Accession Number: BC131624	Purification Method: Antigen affinity purification
Size: 400 µg/ml	GenelD (NCBI): 57645	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	UNIPROT ID: Q9P215	
Isotype: IgG Immunogen Catalog Number: AG21681	Full Name: pogo transposable element with KRAB domain Calculated MW: 609 aa, 69 kDa	
	Tested Applications:	Positive Controls: WB : A549 cells, human testis tissue, MCF-7 cells
Species Specificity: human		
POGK also named as pogo transposable element with KRAB domain is a 609 amino acid protein, which contains of DDE domain, 1 KRAB domain and one HTH CENPB-type DNA binding domain. POGK belongs to DNA-mediated transposons family and may involve in the regulation of gene transcription.		
Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and	·	
	24888-1-AP Size: 400 µ g/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG21681 Tested Applications: WB, ELISA Species Specificity: human POGK also named as pogo transpo DDE domain, 1 KRAB domain and transposons family and may invo Storage: Storage: Storage Buffer:	24888-1-AP BC131624 Size: GeneID (NCBI): 400 μ g/ml 57645 Source: UNIPROT ID: Rabbit Q9P215 Isotype: Full Name: IgG pogo transposable element with Immunogen Catalog Number: AG21681 AG21681 Calculated MW: 609 aa, 69 kDa Observed MW: 85 kDa WB : A545 Species Specificity: WB : A545 human POGK also named as pogo transposable element with KRAB domain is a DDE domain, 1 KRAB domain and one HTH CENPB-type DNA binding doi transposons family and may involve in the regulation of gene transcrip Storage: Storage: Storage Lifter: Storage Buffer:

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



A549 cells were subjected to SDS PAGE followed by western blot with 24888-1-AP (POGK Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.