#### For Research Use Only

# SOX17 Polyclonal antibody

Catalog Number:24903-1-AP

Featured Product

17 Publications

BC111365

GeneID (NCBI):

GenBank Accession Number:



**Purification Method:** 

WB 1:1000-1:4000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: 24903-1-AP Concentration: 500 ug/ml

 500 ug/ml
 64321

 Source:
 UNIPROT ID:

 Rabbit
 Q9H612

 Isotype:
 Full Name:

 IpG
 SRY (sex details)

gG SRY (sex determining region Y)-box

Immunogen Catalog Number:

AG19157 Calculated MW:

414 aa, 44 kDa Observed MW: 44 kDa

**Applications** 

Tested Applications:

WB, ELISA

Cited Applications:

WB, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse, rat Positive Controls:

WB: C6 cells, PC-3 cells, mouse embryo

### **Background Information**

Transcription factor SOX 17 (SOX17) also named as SRY box 17 is a 414 amino acid protein, which contains one Sox C-terminal domain and one HMG box DNA-binding domain. SOX17 as a transcription regulator binds target promoter DNA and bend the DNA via WNT3A. SOX17 is a marker of endodermal cells and a transcriptional regulator containing a DNA binding domain called the HMG box. In mouse embryos, SOX17 plays critical roles in the growth and differentiation of definitive endodermal cells (PMID: 11973269), the remodeling of endothelial cells (PMID: 16895970), hindgut endoderm expansion with localization of primordial germ cells (PMID: 19371732), and gallbladder/bile duct formation (PMID: 19913509).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xiang Xiong	33144185	Cell Signal	WB
Xianmixinuer Abulaiti	28988986	Stem Cell Reports	WB
Jiajia Yang	34034221	Stem Cell Res	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 24903-1-AP (SOX17 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.