For Research Use Only

RNF38 Polyclonal antibody

Catalog Number: 25132-1-AP

1 Publications



Basic Information

Catalog Number:

25132-1-AP

BC033786

Size:

GeneID (NCBI):

700 µg/ml

152006

Source:

Rabbit

Q9H0F5

Isotype:

GeneID (NCBI):

Full Name:

IgG ring finger protein 38
Immunogen Catalog Number: Calculated MW:
AG16054 515 aa, 58 kDa

Observed MW: 43-58 kDa Antigen affinity purification Recommended Dilutions: WB 1:200-1:1000 IHC 1:20-1:200

Purification Method:

IF/ICC 1:50-1:500

Applications

Tested Applications: IF/ICC, IHC, WB,ELISA Cited Applications:

WB, IHC

Species Specificity:

human
Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HepG2 cells,

IHC: human endometrial cancer tissue,

IF/ICC: HepG2 cells,

Background Information

RING finger protein 38 is a protein that in humans is encoded by the RNF38 gene. This gene encodes a protein with a coiled-coil motif and a RING-H2 motif (C3H2C2) at its carboxy-terminus. The RING motif is a zinc-binding domain found in a large set of proteins playing roles in diverse cellular processes including oncogenesis, development, signal transduction, and apoptosis. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. RNF38 is widely expressed with highest levels in testis. RNF38 exists various isoforms and molecular weight of isoforms are 36, 48, 52 and 58 kDa

Notable Publications

Author	Pubmed ID	Journal	Application
Dian Xiong	29581762	J Cancer	WB,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 25132-1-AP (RNF38 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human endometrial cancer tissue slide using 25132-1-AP (RNF38 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human endometrial cancer tissue slide using 25132-1-AP (RNF38 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 25132-1-AP (RNF38 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).